

BOUNDED ULTRAIMAGINARY INDEPENDENCE AND ITS TOTAL MORLEY SEQUENCES

JAMES HANSON

ABSTRACT. We investigate the following model-theoretic independence relation: $b \downarrow_A^{\text{bu}} c$ if and only if $\text{bdd}^{\text{u}}(Ab) \cap \text{bdd}^{\text{u}}(Ac) = \text{bdd}^{\text{u}}(A)$, where $\text{bdd}^{\text{u}}(X)$ is the class of all ultrimaginaries bounded over X . In particular, we sharpen a result of Wagner to show that $b \downarrow_A^{\text{bu}} c$ if and only if $\langle \text{Autf}(\mathbb{M}/Ab) \cup \text{Autf}(\mathbb{M}/Ac) \rangle = \text{Autf}(\mathbb{M}/A)$, and we establish full existence over hyperimaginary parameters (i.e., for any set of hyperimaginaries A and ultrimaginaries b and c , there is a $b' \equiv_A b$ such that $b' \downarrow_A^{\text{bu}} c$). Extension then follows as an immediate corollary.

We also study *total* \downarrow^{bu} -Morley sequences (i.e., A -indiscernible sequences I satisfying $J \downarrow_A^{\text{bu}} K$ for any J and K with $J + K \equiv_A^{\text{EM}} I$), and we prove that an A -indiscernible sequence I is a total \downarrow^{bu} -Morley sequence over A if and only if whenever I and I' have the same Lascar strong type over A , I and I' are related by the transitive, symmetric closure of the relation ‘ $J + K$ is A -indiscernible.’ This is also equivalent to I being ‘based on’ A in a sense defined by Shelah in his early study of simple unstable theories [10].

Finally, we show that for any A and b in any theory T , if there is an Erdős cardinal $\kappa(\alpha)$ with $|Ab| + |T| < \kappa(\alpha)$, then there is a total \downarrow^{bu} -Morley sequence $(b_i)_{i < \omega}$ over A with $b_0 = b$.

INTRODUCTION

A central theme in neostability theory is the importance of various kinds of ‘generic’ indiscernible sequences—usually with Michael Morley’s name attached to them—such as Morley sequences in stable and simple theories, strict Morley sequences in NIP and NTP₂ theories, tree Morley sequences in NSOP₁ theories, and \downarrow^{b} -Morley sequences in rosy theories. A very broad question one might ask is this: How generically can we build indiscernible sequences in *arbitrary* theories?

Over a model M , we can always extend a given type $p(x) \in S_x(M)$ to a global M -invariant type $q(x) \supset p(x)$ and then use this to generate a sequence $(b_i)_{i < \omega}$ satisfying $b_i \models q \upharpoonright M b_{< i}$ for each $i < \omega$. In some cases the particular choice of $q(x)$ matters, but typically these sequences are robustly generic. Sequences produced in this way have a certain property, which is that they are *based on M in the sense of Simon*; i.e., for any I and J with $I \equiv_M J \equiv_M b_{< \omega}$, there is a K such that $I + K$ and $J + K$ are both M -indiscernible. In NIP theories, the sequences with this property are precisely the sequences generated by an invariant type [12, Prop. 2.38]. Over an arbitrary set of parameters A , however, there may fail to be any indiscernible sequences based on A . In the dense circular order, for instance, there are no indiscernible sequences based on \emptyset . Other technical issues arise when

Date: January 10, 2022.

2020 Mathematics Subject Classification. 03C45.

Key words and phrases. ultrimaginaries, bounded closure, total Morley sequences.

working over arbitrary sets, as well, such as the necessity of considering Lascar strong types over and above ordinary types.

A notion of independence \downarrow^* is said to satisfy *full existence* if for any A , b , and c , there is a $b' \equiv_A b$ such that $b' \downarrow_A^* c$. Together with a common model-theoretic application of the Erdős-Rado theorem (Fact 1.2), this implies that for any A and b , one can build an \downarrow^* -Morley sequence, an A -indiscernible sequence $(b_i)_{i < \omega}$ with $b_0 = b$ satisfying $b_i \downarrow_A^* b_{<i}$ for each $i < \omega$ (assuming \downarrow^* also satisfies right monotonicity). Model-theoretically tame theories often have full existence for powerful independence notions, such as non-forking, but this does fail in some notable tame contexts.

One independence notion that is known to satisfy full existence in arbitrary theories is that of *algebraic independence* [2, Prop. 1.5]: $b \downarrow_A^a c$ if $\text{acl}^{\text{eq}}(Ab) \cap \text{acl}^{\text{eq}}(Ac) = \text{acl}^{\text{eq}}(A)$. A natural modification of this concept is *bounded hyperimaginary independence*: $b \downarrow_A^b c$ if $\text{bdd}^{\text{heq}}(Ab) \cap \text{bdd}^{\text{heq}}(Ac) = \text{bdd}^{\text{heq}}(A)$. Despite perhaps sounding like an intro-to-model-theory exercise, the combinatorics necessary to prove full existence for \downarrow^a are somewhat subtle. It was recently established in [5] by Conant and the author that \downarrow^a satisfies full existence in continuous logic and, relatedly, that \downarrow^b satisfies full existence in discrete (and continuous) logic, answering a question of Adler [1, Quest. A.8]. While the relations of \downarrow^a and \downarrow^b are algebraically nice,¹ they seem to lack semantic consequences outside of certain special theories (such as those with a canonical independence relation in the sense of Adler [1, Lem. 3.2]).

While being able to build \downarrow^* -Morley sequences is certainly good, in many applications the important property is really that of being a *total \downarrow^* -Morley sequence*,² which is an A -indiscernible sequence satisfying $b_{\geq i} \downarrow_A^* b_{<i}$ for every $i < \omega$. When \downarrow^* lacks the algebraic properties necessary to imply that all \downarrow^* -Morley sequences are total \downarrow^* -Morley sequences, it can in general be difficult to ensure their existence. Total \downarrow^a -Morley sequences arise in Adler's characterization of canonical independence relations. And building total \downarrow^K -Morley sequences, where \downarrow^K is the relation of non-Kim-forking, is a crucial technical step in Kaplan and Ramsey's proofs of the symmetry of Kim-forking and the independence theorem in NSOP₁ theories [8].

In simple theories, Morley sequences over A are not generally based on A in the sense of Simon. They do however nearly satisfy this property. If I and J are Morley sequences over A with $I \equiv_A^L J$,³ then there are I' and K such that $I + I'$, $I' + K$, and $J + K$ are A -indiscernible. In an NSOP₁ theory T , if I is a tree Morley sequence over $M \models T$ and $J \equiv_M I$, then we can find K_0 , K_1 , and K_2 such that $I + K_0$, $K_1 + K_0$, $K_1 + K_2$, and $J + K_2$ are all M -indiscernible (see Proposition 4.26). This fact suggests the consideration of the following equivalence relation, originally introduced by Shelah in [10, Def. 5.1]: Let \approx_A be the transitive, symmetric closure of the relation ' $I + J$ is A -indiscernible.' The intuition is that what it means for an A -indiscernible sequence I to be 'based on A ' is that there

¹In the sense of the algebra of an independence relation, not the sense of the algebra in 'algebraic closure.'

²This use of the term 'total' in the context of Morley sequences was originally introduced in [8].

³The equivalence relation \equiv_A^L is the transitive closure of the relation 'there is a model $M \supseteq A$ such that $b \equiv_M c$.' If $b \equiv_A^L c$, we say that b and c have the same *Lascar strong type* over A .

are few \approx_A -classes among the realizations of $\text{tp}(I/A)$. We say that I is *based on A in the sense of Shelah* if there does not exist a sequence $(I_i)_{i < \kappa}$ (with κ large) such that $I_i \equiv_A I$ for each $i < \kappa$ and $I_i \not\approx_A I_j$ for each $i < j < \kappa$. A simple compactness argument shows that I is based on A in the sense of Shelah if and only if the set of realizations of $\text{tp}(I/A)$ decomposes into a bounded number of \approx_A -classes. In [4, Def. 2.4],⁴ Buechler used this relation to define a notion of canonical base. He focuses on \emptyset -indiscernible sequences and gives the following definition: A is a *canonical base* of the \emptyset -indiscernible sequence I if any automorphism $\sigma \in \text{Aut}(\mathbb{M})$ fixes A pointwise if and only if it fixes the \approx_\emptyset -class of I . One difficulty with this concept, of course, is that not all indiscernible sequences have canonical bases in this sense (even in T^{eq} , e.g., [1, Ex. 3.13]).

Two of the problems we have mentioned—the lack of canonical bases for indiscernible sequences and the lack of semantic consequences of \downarrow^a and \downarrow^b —can both be solved by an extremely blunt move: the introduction of ultrimaginary parameters. An *ultrimaginary* is an equivalence class of an arbitrary invariant equivalence relation (as opposed to a type-definable equivalence relation, as in the definition of hyperimaginaries). Every indiscernible sequence I trivially has an ultrimaginary canonical base in the sense of Buechler, i.e., the \approx_\emptyset -class of I itself.

Another appealing aspect of ultrimaginaries is that they characterize Lascar strong type in the same way that hyperimaginaries characterize Kim-Pillay strong type. An ultrimaginary $[b]_E$ is *bounded over A* if it has boundedly many conjugates under $\text{Aut}(\mathbb{M}/A)$. We will write $\text{bdd}^u(A)$ for the class of ultrimaginaries bounded over A . In general, it turns out that b and c have the same Lascar strong type over A if and only if they ‘have the same type over $\text{bdd}^u(A)$,’ once this concept is defined precisely.

Pure analogical thinking might lead one to consider the following independence notion: $b \downarrow_A^{\text{bu}} c$ if $\text{bdd}^u(Ab) \cap \text{bdd}^u(Ac) = \text{bdd}^u(A)$. This notion is implicit in a result of Wagner [13, Prop. 2.12], which we restate and expand slightly (Proposition 2.4): $b \downarrow_A^{\text{bu}} c$ if and only if $\langle \text{Autf}(\mathbb{M}/Ab) \cup \text{Autf}(\mathbb{M}/Ac) \rangle = \text{Autf}(\mathbb{M}/A)$ (where $\langle X \rangle$ is the group generated by X). This characterization is clearly semantically meaningful, and moreover it allows one to discuss \downarrow^{bu} without actually mentioning ultrimaginaries at all. One way to see why this equivalence works is the fact that ultrimaginaries are ‘dual’ to co-small sets of automorphisms; a group $G \leq \text{Aut}(\mathbb{M})$ is *co-small* if there is a small model M such that $\text{Aut}(\mathbb{M}/M) \leq G$. For every co-small group G , there is an ultrimaginary a_E such that $\text{Aut}(\mathbb{M}/a_E) = G$ (Proposition 1.7).

As \downarrow^{bu} lacks finite character, *total \downarrow^{bu} -Morley sequences over A* seem to be correctly defined as A -indiscernible sequences $(b_i)_{i < \omega}$ with the property that for any $I + J \equiv_A^{\text{EM}} b_{< \omega}$,⁵ we have that $I \downarrow_A^{\text{bu}} J$. The automorphism group characterization of \downarrow^{bu} , together with its nice algebraic properties and the malleability of indiscernible sequences, leads to a pleasing characterization of total \downarrow^{bu} -Morley

⁴This preprint is difficult to track down. The relevant ideas are developed further by Adler in [1, Sec. 3.2], which is easily available.

⁵ $I \equiv_A^{\text{EM}} J$ means that I and J have the same *Ehrenfeucht-Mostowski type* over A (i.e., for any increasing tuples $\bar{b} \in I$ and $\bar{c} \in J$ of the same length, $\bar{b} \equiv_A \bar{c}$). Note that I and J do not need to have the same order type.

sequences over sets of hyperimaginary parameters (Theorem 4.8), the equivalence of the following.

- $(b_i)_{i < \omega}$ is a total \downarrow^{bu} -Morley sequence over A .
- For some infinite I and J , we have that $I + J \equiv_A^{\text{EM}} b_{<\omega}$ and $I \downarrow_A^{\text{bu}} J$.
- For any I , $I \approx_A b_{<\omega}$ if and only if there is $I' \equiv_A^{\text{L}} I$ such that $b_{<\omega} + I'$ is A -indiscernible.
- $b_{<\omega}$ is based on A in the sense of Shelah; i.e., $[b_{<\omega}]_{\approx_A} \in \text{bdd}^{\text{u}}(A)$.

The condition in the third bullet point is a natural mutual generalization of Lascar strong type and Ehrenfeucht-Mostowski type (Definition 4.5). Theorem 4.8 also tells us that when total \downarrow^{bu} -Morley sequences exist, they act as particularly uniform witnesses of Lascar strong type (Proposition 4.3).

Of course this all leave two critical questions: Does \downarrow^{bu} always satisfy full existence? And, even if it does, can we actually build total \downarrow^{bu} -Morley sequences in any type over any set under any theory? The bluntness of ultraimaginaries leaves us without one of the most important tools in model theory, compactness. Furthermore, \downarrow^{bu} 's lack of finite character gives us less leeway in applying the Erdős-Rado theorem to construct indiscernible sequences with certain properties; we now need to be more concerned with the particular order types of the sequences involved.

Using some of the indiscernible tree technology from [8], we are able to prove that \downarrow^{bu} does satisfy full existence over arbitrary sets of (hyperimaginary) parameters in arbitrary (discrete or continuous) theories (Theorem 3.5).⁶ With regards to building total \downarrow^{bu} -Morley sequences, Theorem 4.8 tells us that we don't need to worry *too* much about order types. All we need to get a total \downarrow^{bu} -Morley sequence over A is an A -indiscernible sequence $(b_i)_{i < \omega + \omega}$ with $b_{\geq \omega} \downarrow_A^{\text{bu}} b_{<\omega}$. This is fortunate because constructing ill-ordered \downarrow^{bu} -Morley sequences directly seems daunting. Unfortunately, $\omega + \omega$ appears to be about one ω further than we can go without a large cardinal. What we do get is this (Theorem 4.21): For any A and b in any theory T , if there is an Erdős cardinal $\kappa(\alpha)$ with $|Ab| + |T| < \kappa(\alpha)$ (for any $\alpha \geq \omega$), then there is a total \downarrow^{bu} -Morley sequence $(b_i)_{i < \omega}$ over A with $b_0 = b$. Without a large cardinal, the best we seem to be able to do (Proposition 4.16) is a half-infinite, half-arbitrary-finite approximation of a total \downarrow^{bu} -Morley sequence, which we call a *weakly total \downarrow^{bu} -Morley sequence*. These sequences also serve as uniform witnesses of Lascar strong type, without any set-theoretic hypotheses (Corollary 4.17).

1. ULTRAIMAGINARIES

Here we will set definitions and conventions, and we also take the opportunity to collect some basic facts about ultraimaginaries which are likely folklore, although we could not find explicit references.

Fix a theory T and a monster model $\mathbb{M} \models T$.

Definition 1.1. An *invariant equivalence relation of arity κ* is an equivalence relation E on \mathbb{M}^x (with $|x| = \kappa$) such that for any $a, b, c, d \in \mathbb{M}^x$ with $ab \equiv cd$, aEb if and only if cEd .

⁶Although this result partially supersedes a result in [5] (full existence for \downarrow^{a} in continuous logic and \downarrow^{b} in discrete or continuous logic), the proof there gives more detailed numerical information which may be especially useful in the metric context.

An *ultraimaginary of arity κ* is an E -equivalence class a_E of some tuple $a \in \mathbb{M}^x$ under an invariant equivalence relation E of arity κ . We may also write such an equivalence class as $[a]_E$ if necessary for notational clarity.

Given an ultraimaginary a_E , $\text{Aut}(\mathbb{M}/a_E)$ is the set of automorphisms $\sigma \in \text{Aut}(\mathbb{M})$ with the property that $aE(\sigma \cdot a)$. We write $\text{Autf}(\mathbb{M}/a_E)$ for the group generated by $\{\sigma \in \text{Aut}(\mathbb{M}/M) : M \preceq \mathbb{M}, \text{Aut}(\mathbb{M}/M) \leq \text{Aut}(\mathbb{M}/a_E)\}$.

We say that b_F is *definable over a_E* if b_F is fixed by every automorphism of $\text{Aut}(\mathbb{M}/a_E)$. We write $\text{dcl}^u(a_E)$ for the class of all ultraimaginaries definable over a_E . For any κ , we write $\text{dcl}_\kappa^u(a_E)$ for the set of elements of $\text{dcl}^u(a_E)$ of arity at most κ . We say that b_F and c_G are *interdefinable over a_E* if $b_F \in \text{dcl}^u(a_E c_G)$ and $c_G \in \text{dcl}^u(a_E b_F)$.

We say that b_F is *bounded over a_E* if the $\text{Aut}(\mathbb{M}/b_F)$ -orbit of a_E is bounded. We write $\text{bdd}^u(a_E)$ for the class of all ultraimaginaries bounded over a_E . We write $\text{bdd}_\kappa^u(a_E)$ for the set of elements of $\text{bdd}^u(a_E)$ of arity at most κ . We say that b_F and c_G are *interbounded over a_E* if $b_F \in \text{bdd}^u(a_E c_G)$ and $c_G \in \text{bdd}^u(a_E b_F)$.

We write $a_E \equiv b_E$ to mean that there is an automorphism $\sigma \in \text{Aut}(\mathbb{M})$ with $\sigma \cdot a_E = b_E$. We write $b_F \equiv_{a_E} c_F$ to mean that $a_E b_F \equiv a_E c_F$ (i.e., there is $\sigma \in \text{Aut}(\mathbb{M}/a_E)$ such that $\sigma \cdot b_F = c_F$).

We will also sometimes define an invariant equivalence relation E on the realizations of a single type $p(x)$ over \emptyset . Equivalence classes of such can be thought of as ultraimaginaries by using the same trick that is commonly used with hyperimaginaries: Consider the invariant equivalence relation $E'(x, y)$ defined by $(x = y \wedge x \not\models p) \vee (E(x, y) \wedge x \models p \wedge y \models p)$.

For the sake of clarity, we will reserve the notation a_E for ultraimaginaries and write hyperimaginaries in the same way we write real elements. For the sake of cardinality issues, we will also take all hyperimaginaries to be quotients of countable tuples by countably type-definable equivalence relations. It is a standard fact that every hyperimaginary is interdefinable with some set of hyperimaginaries of this form. Both of these conventions can be justified more rigorously by passing to the ‘continuous version of T^{eq} .’ See the discussion in [5, Sec. 4.3] for details.

Fact 1.2 ([3, Lem. 1.2]). *Let $(b_i)_{i < \lambda}$ be a sequence of tuples with $|b_i| < \kappa$ and let A be some set of parameters. If $\lambda \geq \beth_{(2^\kappa + |A| + |T|)^+}$, then there is an A -indiscernible sequence $(b'_i)_{i < \omega}$ such that for every $n < \omega$, there are $i_0 < \dots < i_n < \kappa$ such that $b'_0 \dots b'_n \equiv_A b_{i_0} \dots b_{i_n}$.*

Lemma 1.3. *Let M be a model. If $a_E \in \text{bdd}^u(M)$, then $a_E \in \text{dcl}^u(M)$.*

Proof. Assume that $a_E \notin \text{dcl}^u(M)$. Let $p(x)$ be a global M -invariant type extending $\text{tp}(a/M)$. Assume that there are a_0 and a_1 realizing $\text{tp}(a/M)$ such that $a_0 \not\equiv a_1$. For any $i > 1$, given $a_{<i}$, let $a_i \models p \upharpoonright M a_{<i}$. Since $a_i a_j \equiv_M a_i a_k$ for any $j, k < i$, we must have that $a_i \not\equiv a_j$ for any $j < i$. Since we can do this indefinitely, we have that a_E is not bounded over M . \square

Proposition 1.4. *For any ultraimaginaries a_E and b_F , the following are equivalent.*

- (1) $b_F \notin \text{bdd}^u(a_E)$.
- (2) There is an a -indiscernible sequence $(b_i)_{i < \omega}$ such that $b_0 \equiv_{a_E} b$ and $b_i \not\equiv b_j$ for each $i < j < \omega$.
- (3) $|\text{Aut}(\mathbb{M}/a_E) \cdot b_F| > 2^{|ab| + |T|}$.

Proof. (1) \Rightarrow (2). This follows from an immediate application of Fact 1.2.

(2) \Rightarrow (3). This follows by stretching the relevant indiscernible sequence to a length of $(2^{|ab|+|T|})^+$.

(3) \Rightarrow (1). Let $(b_F^i)_{i < (2^{|ab|+|T|})^+}$ be an enumeration of $\text{Aut}(\mathbb{M}/a_E) \cdot b_F$. Let $M \supseteq a$ be a model with $|M| \leq |a| + |T|$. Let x be a tuple of variables of the same length as b . There are at most $2^{|ab|+|T|}$ types in $S_x(M)$. Therefore, there must be $i < j < (2^{|ab|+|T|})^+$ such that $b^i \equiv_M b^j$. Let $p(x)$ be a global M -invariant type extending $\text{tp}(b^i/M)$, and let $(c^i)_{i < \lambda}$ be a Morley sequence generated by $p(x)$ over $Mb^i b^j$. Since $b^i \not\equiv b^j$, we must have that $c^0 \not\equiv b^i$. Therefore $c^i \not\equiv c^j$ for any $i < j < \lambda$. Furthermore, since $c^i \equiv_M c^j$, we also have that $c^i \equiv_a c^j$ and therefore also $c_F^i \equiv_{a_E} c_F^j \equiv_{a_E} b_F$. Since we can do this for any λ , we have that $b_F \notin \text{bdd}^u(a_E)$. \square

Corollary 1.5. *For any λ , $\text{bdd}_\lambda^u(a_E)$ has cardinality at most $2^{|a|+2^{\lambda+|T|}}$.*

Proof. There are at most $2^{2^{\lambda+|T|}}$ many invariant equivalence relations of arity at most λ . For each such F , the set $\{b_F : b_F \in \text{bdd}_\lambda^u(a_E)\}$ has cardinality at most $2^{|a|+\lambda+|T|}$ by Proposition 1.4. Finally, $2^{2^{\lambda+|T|}} \cdot 2^{|a|+\lambda+|T|} = 2^{|a|+2^{\lambda+|T|}}$. \square

1.1. Co-small groups of automorphisms. Here we will see that ultramaginarities are essentially the same thing as reasonable subgroups of $\text{Aut}(\mathbb{M})$.

Definition 1.6. A set $G \leq \text{Aut}(\mathbb{M})$ is *co-small* if there is a small model M such that $\text{Aut}(\mathbb{M}/M) \leq G$.

Clearly for any ultramaginary a_E , $\text{Aut}(\mathbb{M}/a_E)$ is co-small. The converse is true as well.

Proposition 1.7. *For any co-small G , if $\text{Aut}(\mathbb{M}/M) \leq G$, then there is an ultramaginary a_E such that $G = \text{Aut}(\mathbb{M}/a_E)$ where a is some enumeration of M .*

Proof. Let M be a small model witnessing that G is co-small. Consider the binary relation defined on realizations of $\text{tp}(M)$ (in some fixed enumeration) defined by $E(M_0, M_1)$ if and only if there is $\sigma \in \text{Aut}(\mathbb{M})$ and $\tau \in G$ such that $\sigma \cdot M = M_0$ and $\sigma\tau \cdot M = M_1$. We need to verify that E is an invariant equivalence relation. Reflexivity is obvious.

Invariance. Suppose that $E(M_0, M_1)$, as witnessed by $\sigma \in \text{Aut}(\mathbb{M})$ and $\tau \in G$. Fix $\sigma' \in \text{Aut}(\mathbb{M})$. We then have that $\sigma'\sigma \cdot M = \sigma' \cdot M_0$ and $\sigma'\sigma\tau \cdot M = \sigma' \cdot M_1$, whence $E(\sigma' \cdot M_0, \sigma' \cdot M_1)$.

Symmetry. If $\sigma \cdot M = M_0$ and $\sigma\tau \cdot M = M_1$ with $\sigma \in \text{Aut}(\mathbb{M})$ and $\tau \in G$, then $\sigma\tau\tau^{-1} \cdot M = M_0$ and $\sigma\tau \cdot M = M_1$. We have $\sigma\tau \in \text{Aut}(\mathbb{M})$ and $\tau^{-1} \in G$, so $E(M_1, M_0)$.

Transitivity. Suppose that for $\sigma, \sigma' \in \text{Aut}(\mathbb{M})$ and $\tau, \tau' \in G$, we have that $\sigma \cdot M = M_0$, $\sigma\tau \cdot M = \sigma' \cdot M = M_1$, and $\sigma'\tau' \cdot M = M_2$. This implies that $(\sigma\tau)^{-1}\sigma' = \tau^{-1}\sigma^{-1}\sigma' \in \text{Aut}(\mathbb{M}/M) \leq G$. Since $\tau \in G$ as well, we have that $\sigma^{-1}\sigma' \in G$. Therefore $\sigma^{-1}\sigma'\tau' \in G$. Finally, $\sigma\sigma^{-1}\sigma'\tau' \cdot M = M_2$, so $E(M_0, M_2)$.

Consider the ultramaginary M_E . For any $\tau \in G$, we clearly have $E(M, \tau \cdot M)$, so $G \leq \text{Aut}(\mathbb{M}/M_E)$. Conversely, suppose that $\alpha \in \text{Aut}(\mathbb{M}/M_E)$. By definition, this implies that $E(M, \alpha \cdot M)$, so there are $\sigma \in \text{Aut}(\mathbb{M})$ and $\tau \in G$ such that $\sigma \cdot M = M$ and $\sigma\tau \cdot M = \alpha \cdot M$. Therefore $\sigma, \tau^{-1}\sigma^{-1}\alpha \in \text{Aut}(\mathbb{M}/M) \leq G$. Since $\tau^{-1} \in G$, we therefore have that $\alpha \in G$. \square

Corollary 1.8. *If $b_F \in \text{bdd}^u(a_E)$, then there is $c_G \in \text{bdd}^u(a_E)$ of arity at most $|a| + |T|$ such that b_F and c_G are interdefinable over \emptyset . Furthermore, c can be taken to be an enumeration of any model of size at most $|a| + |T|$ containing a .*

Proof. There is a model $M \supseteq a$ with $|M| \leq |a| + |T|$. By Lemma 1.3, we have that $\text{Aut}(M/M) \leq \text{Aut}(M/b_F)$, so by Proposition 1.7, we have that there is c_G with arity at most $|a| + |T|$ which satisfies that $\text{Aut}(M/c_G) = \text{Aut}(M/b_F)$ (i.e., c_G and b_F are interdefinable over \emptyset). Furthermore, we can take c to be an enumeration of M . \square

Definition 1.9. For any co-small group G , we write $[[G]]$ for some arbitrary ultra-imaginary a_E of minimal arity satisfying $G = \text{Aut}(M/a_E)$. We will write $\text{dcl}^u[[G]]$ and $\text{dcl}_\lambda^u[[G]]$ for $\text{dcl}^u([[G]])$ and $\text{dcl}_\lambda^u([[G]])$ and likewise with bdd^u . (Note that $\text{dcl}^u[[G]]$ and $\text{bdd}^u[[G]]$ only depend on G , not on the particular choice of $[[G]]$.)

It is immediate from Proposition 1.7 that for any co-small G and H , $[[G]] \in \text{dcl}^u[[H]]$ if and only if $G \geq H$. A similar statement is true of bdd^u (Proposition 1.12).

Now we can see that intersections of dcl^u -closed sets (and therefore also bdd^u -closed sets) have semantic significance in arbitrary theories, in that intersections correspond to joins in the lattice of co-small groups of automorphisms.

Proposition 1.10. *For any a_E, b_F, c_G , and c'_G , the following are equivalent.*

- (1) $c_G \equiv_{\text{dcl}_\lambda^u(a_E) \cap \text{dcl}_\lambda^u(b_F)} c'_G$ for all λ .
- (2) There is $\sigma \in \langle \text{Aut}(M/a_E) \cup \text{Aut}(M/b_F) \rangle$ such that $\sigma \cdot c_G = c'_G$.
- (3) There is a sequence $(a^i b^i c^i)_{i \leq n}$ such that $a^0 = a, b^0 = b, c^0 = c, c^n = c'$, and for each $i < n$,
 - if i is even, then $a^i = a^{i+1}$ and $b_F^i c_G^i \equiv_{a_E^i} b_F^{i+1} c_G^{i+1}$ and
 - if i is odd, then $b^i = b^{i+1}$ and $a_E^i c_G^i \equiv_{b_F^i} a_E^{i+1} c_G^{i+1}$.

Proof. Let $H = \langle \text{Aut}(M/a_E) \cup \text{Aut}(M/b_F) \rangle$.

Claim. $\text{dcl}_\lambda^u(a_E) \cap \text{dcl}_\lambda^u(b_F)$ and $[[H]]$ are interdefinable for sufficiently large λ .

Proof of claim. Clearly $[[H]] \in \text{dcl}^u(a_E) \cap \text{dcl}^u(b_F)$, so $[[H]] \in \text{dcl}_\lambda^u(a_E) \cap \text{dcl}_\lambda^u(b_F)$ for all sufficiently large λ .

Conversely, suppose that $d_I \in \text{dcl}^u(a_E) \cap \text{dcl}^u(b_F)$. Any $\sigma \in H$ is a product of elements of $\text{Aut}(M/a_E)$ and $\text{Aut}(M/b_F)$, so it must fix d_I . Therefore $\text{Aut}(M/d_I) \geq H$ and hence $d_I \in \text{dcl}^u[[H]]$. \square_{claim}

So now we have that $c_G \equiv_{\text{dcl}_\lambda^u(a_E) \cap \text{dcl}_\lambda^u(b_F)} c'_G$ holds for sufficiently large λ if and only if $c_G \equiv_{[[H]]} c'_G$. Also note that $c_G \equiv_{\text{dcl}_\lambda^u(a_E) \cap \text{dcl}_\lambda^u(b_F)} c'_G$ for sufficiently large λ and only if the same holds for any λ . Therefore we have that (1) and (2) are equivalent.

There is a $\sigma \in H$ with $\sigma \cdot c_G = c'_G$ if and only if there are $\alpha_0, \dots, \alpha_n \in \text{Aut}(M/a_E)$ and $\beta_0, \dots, \beta_n \in \text{Aut}(M/b_E)$ such that $\sigma = \beta_n \alpha_n \beta_{n-1} \dots \alpha_1 \beta_0 \alpha_1$. The existence of such $\bar{\alpha}$ and $\bar{\beta}$ for which $\beta_n \alpha_n \beta_{n-1} \dots \alpha_1 \beta_0 \alpha_1 \cdot c_G = c'_G$ is clearly equivalent to (3), so we have that (2) and (3) are equivalent. \square

A similar statement is true for intersections of arbitrary families of small sets.

1.2. Lascar strong type.

Definition 1.11. For any co-small group $G \leq \text{Aut}(\mathbb{M})$, let G_f be the group generated by all groups of the form $\text{Aut}(\mathbb{M}/M) \leq G$ with M a small model. For any ultrimagininary a_E , let $\text{Autf}(\mathbb{M}/a_E) = \text{Aut}(\mathbb{M}/a_E)_f$.

We say that b_F and c_F have the same *Lascar strong type over a_E* , written $b_F \equiv_{a_E}^L c_F$, if there is $\sigma \in \text{Autf}(\mathbb{M}/a_E)$ such that $\sigma \cdot b_F = c_F$.

Proposition 1.12. *For any co-small groups G and H , $[[G]] \in \text{bdd}^u[[H]]$ if and only if $G \geq H_f$.*

Proof. First note that for a model M , by Lemma 1.3, we have that $[[G]] \in \text{bdd}^u(M)$ if and only if $G \geq \text{Aut}(\mathbb{M}/M)$. Therefore, for any model M with $[[H]] \in \text{bdd}^u(M)$, we must have that $G \geq \text{Aut}(\mathbb{M}/M)$. Since $[[H]] \in \text{bdd}^u(M)$ if and only if $H \geq \text{Aut}(\mathbb{M}/M)$, we have that $G \geq H_f$.

Conversely, assume that $G \geq H_f$. This implies that for any small model M with $[[H]] \in \text{bdd}^u(M)$, we have $H_f \geq \text{Aut}(\mathbb{M}/M)$, so $G \geq \text{Aut}(\mathbb{M}/M)$ and $[[G]] \in \text{dcl}^u(M)$. Fix some such model N . Assume for the sake of contradiction that $[[G]] \notin \text{bdd}^u[[H]]$. For any λ , we can find $(\sigma_i)_{i < \lambda}$ in $H = \text{Aut}(\mathbb{M}/[[H]])$ such that $\sigma_i \cdot [[G]] \neq \sigma_j \cdot [[G]]$ for each $i < j < \lambda$. If λ is larger than $2^{|N|+|T|}$, there must be $i < j < \lambda$ such that $\sigma_i \cdot [[G]] \equiv_N \sigma_j \cdot [[G]]$. Let $N' = \sigma_i^{-1} \cdot N$. N' is now a model satisfying $\text{Aut}(\mathbb{M}/N') \leq G$. So $[[G]] \in \text{dcl}^u(N')$, but $[[G]] \equiv_{N'} \sigma_i^{-1} \sigma_j \cdot [[G]]$ and $[[G]] \neq \sigma_i^{-1} \sigma_j \cdot [[G]]$, which is a contradiction. \square

An important fact about ultrimaginaries is that bdd^u has the same relationship with Lascar strong type that bdd^{heq} has with Kim-Pillay strong type.

For any a_E and b_F , by an abuse of notation, we'll write $[b_F]_{\equiv_{a_E}^L}$ for $[ab]_G$, where $G(ab, a'c')$ holds if and only if aEa' and $b \equiv_{a_E}^L b'$.

Proposition 1.13. *For any ultrimaginaries a_E , b_F , and c_F , the following are equivalent.*

- (1) $b_F \equiv_{a_E}^L c_F$.
- (2) $b_F \equiv_{\text{bdd}_\lambda^u(a_E)} c_F$ for all sufficiently large λ .
- (3) $b_F \equiv_{\text{bdd}_{|a|+|T|}^u(a_E)} c_F$.

Proof. To see that (1) implies (3), fix $M \supseteq a$ and some automorphism $\sigma \in \text{Aut}(\mathbb{M}/M)$. By Lemma 1.3, we have that $\text{Aut}(\mathbb{M}/M) \leq \text{Aut}(\mathbb{M}/\text{bdd}_{|a|+|T|}^u(a_E))$. Therefore $b_F \equiv_{\text{bdd}_{|a|+|T|}^u(a_E)} \sigma \cdot b_F$. By induction, we therefore have that $b_F \equiv_{a_E}^L c_F$ implies $b_F \equiv_{\text{bdd}_{|a|+|T|}^u(a_E)} c_F$.

Corollary 1.8 implies that $\text{Aut}(\mathbb{M}/\text{bdd}_\lambda^u(a_E)) \geq \text{Aut}(\mathbb{M}/\text{bdd}_{|a|+|T|}^u(a_E))$ for all λ , so (3) implies (2).

To see that (2) implies (1), note that $[b_F]_{\equiv_{a_E}^L} \in \text{bdd}_\lambda^u(a_E)$ for some sufficiently large λ . Therefore if $b_F \equiv_{\text{bdd}_\lambda^u(a_E)} c_F$, we must have $[b_F]_{\equiv_{a_E}^L} = [c_F]_{\equiv_{a_E}^L}$ or, in other words, $b_F \equiv_{a_E}^L c_F$. \square

2. BOUNDED ULTRAIMAGINARY INDEPENDENCE

Definition 2.1. We write $b_F \downarrow_{a_E}^{\text{bu}} c_G$ to mean that $\text{bdd}^u(a_E b_F) \cap \text{bdd}^u(a_E c_G) = \text{bdd}^u(a_E)$.

An easy argument shows that if $c_G \in \text{bdd}^u(b_F)$ and $b_F \in \text{bdd}^u(a_E)$, then $c_G \in \text{bdd}^u(a_E)$. Note also that $b_F \downarrow_{a_E}^{\text{bu}} c_G$ if and only if $\text{bdd}_\kappa^u(a_E b_F) \cap \text{bdd}_\kappa^u(a_E c_G) = \text{bdd}_\kappa^u(a_E)$ for all κ . \downarrow^{bu} satisfies some of the familiar properties of \downarrow^{a} .

Proposition 2.2. *Fix ultraimaginararies a_E, b_F, c_G , and e_I .*

- (Invariance) *If $a_E b_F c_G \equiv a'_E b'_F c'_G$, then $b_F \downarrow_{a_E}^{\text{bu}} c_G$ if and only if $b'_F \downarrow_{a'_E}^{\text{bu}} c'_G$.*
- (Symmetry) *$b_F \downarrow_{a_E}^{\text{bu}} c_G$ if and only if $c_G \downarrow_{a_E}^{\text{bu}} b_F$.*
- (Monotonicity) *If $b_F c_G \downarrow_{a_E}^{\text{bu}} d_H e_I$, then $b_F \downarrow_{a_E}^{\text{bu}} d_H$.*
- (Transitivity) *If $b_F \downarrow_{a_E}^{\text{bu}} c_G$ and $d_H \downarrow_{a_E b_F}^{\text{bu}} c_G$, then $b_F d_H \downarrow_{a_E}^{\text{bu}} c_G$.*
- (Normality) *If $b_F \downarrow_{a_E}^{\text{bu}} c_G$, then $a_E b_F \downarrow_{a_E}^{\text{bu}} a_E c_G$.*
- (Anti-reflexivity) *If $b_F \downarrow_{a_E}^{\text{bu}} b_F$, then $b_F \in \text{bdd}^{\text{u}}(a_E)$.*

Proof. Everything except transitivity is immediate. For transitivity, assume that $b_F \downarrow_{a_E}^{\text{bu}} c_G$ and $d_H \downarrow_{a_E b_F}^{\text{bu}} c_G$. Let e_I be an element of $\text{bdd}^{\text{u}}(a_E b_F d_H) \cap \text{bdd}^{\text{u}}(a_E c_G)$. This implies that it is an element of $\text{bdd}^{\text{u}}(a_E b_F d_H) \cap \text{bdd}^{\text{u}}(a_E b_F c_G)$, so by assumption it is an element of $\text{bdd}^{\text{u}}(a_E b_F)$. But this means that it's in both $\text{bdd}^{\text{u}}(a_E b_F)$ and $\text{bdd}^{\text{u}}(a_E c_G)$, so by assumption again, it is an element of $\text{bdd}^{\text{u}}(a_E)$. \square

Part of the goal of this paper is to prove full existence and therefore also extension for \downarrow^{bu} (although only over hyperimaginary bases).

- (Full existence over hyperimaginaries) For any set of hyperimaginaries A and ultraimaginararies b_E and c_F , there is $c'_F \equiv_A c_F$ such that $b_E \downarrow_A^{\text{bu}} c'_F$.
- (Extension over hyperimaginaries) For any set of hyperimaginaries A and ultraimaginararies b_E, c_F , and d_G , if $b_E \downarrow_A^{\text{bu}} c_F$, then there is $b'_E \equiv_{A c_F} b_E$ such that $b'_E \downarrow_A^{\text{bu}} c_F d_G$.

Finite character fails very badly, of course. Local character seems unlikely except possibly in the presence of large cardinals. We do have some control over the relevant cardinalities, however.

Proposition 2.3. *For any a_E, b_F , and c_G , $b_F \downarrow_{a_E}^{\text{bu}} c_G$ if and only if $\text{bdd}^{\text{u}}_{\lambda}(a_E b_F) \cap \text{bdd}^{\text{u}}_{\lambda}(a_E c_G) = \text{bdd}^{\text{u}}_{\lambda}(a_E)$, where $\lambda = |ab| + |T|$.*

Proof. Let $\lambda = |ab| + |T|$. Clearly we have that if $b_F \downarrow_{a_E}^{\text{bu}} c_G$, then $\text{bdd}^{\text{u}}_{\lambda}(a_E b_F) \cap \text{bdd}^{\text{u}}_{\lambda}(a_E c_G) = \text{bdd}^{\text{u}}_{\lambda}(a_E)$.

Conversely, assume that $b_F \not\downarrow_{a_E}^{\text{bu}} c_G$. There is some $d_H \in (\text{bdd}^{\text{u}}(a_E b_F) \cap \text{bdd}^{\text{u}}(a_E c_G)) \setminus \text{bdd}^{\text{u}}(a_E)$. By Corollary 1.8, there is e_I of arity at most λ such that d_H and e_I are interdefinable. This means that $e_I \in (\text{bdd}^{\text{u}}_{\lambda}(a_E b_F) \cap \text{bdd}^{\text{u}}_{\lambda}(a_E c_G)) \setminus \text{bdd}^{\text{u}}_{\lambda}(a_E)$. Therefore $\text{bdd}^{\text{u}}_{\lambda}(a_E b_F) \cap \text{bdd}^{\text{u}}_{\lambda}(a_E c_G) \neq \text{bdd}^{\text{u}}_{\lambda}(a_E)$. \square

The following characterization of \downarrow^{bu} (and the manner of proof) is essentially due to Wagner [13].

Proposition 2.4. *For any ultraimaginararies a_E, b_F , and c_G , the following are equivalent.*

- (1) $b_F \downarrow_{a_E}^{\text{bu}} c_G$.
- (2) For any $b'_F \equiv_{a_E}^{\text{L}} b_F$, there are $b^0, c^0, b^1, c^1, \dots, c^{n-1}, b^n$ such that $b^0 = b$, $c^0 = c$, $b^n = b'$, and for each $i < n$, $b_F^i \equiv_{a_E c_G}^{\text{L}} b_F^{i+1}$ and $c_G^i \equiv_{a_E b_F^{i+1}} c_G^{i+1}$ if $i < n-1$.
- (3) $\langle \text{Autf}(\mathbb{M}/a_E b_F) \cup \text{Autf}(\mathbb{M}/a_E c_G) \rangle = \text{Autf}(\mathbb{M}/a_E)$.

Proof. The equivalence of (1) and (3) follows from Propositions 1.10 and 1.13 and the fact that intersections of bdd^u -closed sets are bdd^u -closed. These propositions also clearly give that (3) implies (2) (after some re-indexing of the relevant sequences).

So assume (2), but also assume for the sake of contradiction that (1) fails. Let d_H be an element of $(\text{bdd}^u(a_E b_F) \cap \text{bdd}^u(a_E c_G)) \setminus \text{bdd}^u(a_E)$. Since d_H is not bounded over a_E , there must be some $d'_H \equiv_{a_E}^L d_H$ such that $d'_H \notin \text{bdd}^u(a_E b_E) \cap \text{bdd}^u(a_E c_G)$. Find b'_F such that $b_F d_H \equiv_{a_E}^L b'_F d'_H$. Let $b^0, c^0, b^1, c^1, \dots, c^{n-1}, b^n$ be as in (2), with $b^n = b'$. Find $d^{1/2}, d^1, d^{3/2}, d^2, \dots, d^{n-1/2}, d^n$ such that $d^{1/2} = d$ and for each $i < n$,

- $b_F^i d_H^{i+1/2} \equiv_{a_E c_G^i}^L b_F^{i+1} d^{i+1}$ and
- $c_G^i d_H^{i+1} \equiv_{a_E b_F^{i+1}}^L c_G^{i+1} d_H^{i+3/2}$ if $i < n-1$.

We now have that $b'_F d'_H \equiv_{a_E}^L b_F d_H \equiv_{a_E}^L b'_F d'_H$, so in particular, $d'_H \equiv_{a_E b'_F}^L d_H^n$. But note that for each $i < n$, we have that

$$\text{bdd}^u(a_E b_F^i) \cap \text{bdd}^u(a_E c_G^i) = \text{bdd}^u(a_E b_F^{i+1}) \cap \text{bdd}^u(a_E c_G^i)$$

and, if $i < n-1$,

$$\text{bdd}^u(a_E b_F^{i+1}) \cap \text{bdd}^u(a_E c_G^i) = \text{bdd}^u(a_E b_F^{i+1}) \cap \text{bdd}^u(a_E c_G^{i+1}).$$

Therefore $d_H^n \in \text{bdd}^u(a_E b_F^n) \cap \text{bdd}^u(a_E c_G^{n-1})$, so since $d_H^n \equiv_{a_E b_F^n}^L d'_H$, we must also have $d'_H \in \text{bdd}^u(a_E b_F^n) \cap \text{bdd}^u(a_E c_G^{n-1}) = \text{bdd}^u(a_E b_F) \cap \text{bdd}^u(a_E c_G)$, which is a contradiction. \square

3. FULL EXISTENCE

We will use the tree bookkeeping machinery from [8], with some minor extensions (the notation \mathcal{T}_α^* and \mathcal{F}_α).

Definition 3.1. For any ordinal α , $\mathcal{L}_{s,\alpha}$ is the language

$$\{\leq, \wedge, <_{lex}, P_0, P_1, \dots, P_\beta(\beta < \alpha), \dots\},$$

with \leq and $<_{lex}$ binary relations, \wedge a binary function, and each P_β a unary relation.

For any ordinal α , we write \mathcal{T}_α^* for the set of functions f with codomain ω and finite support such that $\text{dom}(f)$ is an end segment of α . We write \mathcal{T}_α for the set of functions $f \in \mathcal{T}_\alpha^*$ with $\text{dom}(f) = [\beta, \alpha)$ and β is not a limit ordinal. We write $\mathcal{F}_{\alpha+1}$ (for forest) for $\mathcal{T}_{\alpha+1} \setminus \{\emptyset\}$.

We interpret \mathcal{T}_α^* and \mathcal{T}_α as $\mathcal{L}_{s,\alpha}$ -structures by

- $f \leq g$ if and only if $f \subseteq g$;
- $f \wedge g = f \upharpoonright [\beta, \alpha) = g \upharpoonright [\beta, \alpha)$, where $\beta = \min\{\gamma : f \upharpoonright [\gamma, \alpha) = g \upharpoonright [\gamma, \alpha)\}$ (with the understanding that $\min \emptyset = \alpha$);
- $f <_{lex} g$ if and only if either $f \triangleleft g$ or f and g are \leq -incomparable, $\text{dom}(f \wedge g) = [\gamma, \alpha)$, and $f(\gamma) < g(\gamma)$; and
- $P_\beta(f)$ holds if and only if $\text{dom}(f) = [\beta, \alpha)$.

We write $\langle i \rangle$ for the element $\{(\alpha, i)\}$ of \mathcal{T}_α^* . For $f \in \mathcal{T}_\alpha^*$ with $\text{dom}(f) = [\beta+1, \alpha)$ and $i < \delta$, we write $f \frown \langle i \rangle$ for the function $f \cup \{(\beta, i)\}$. In general, we write $\langle i \rangle \frown f$ for the element of $\mathcal{T}_{\alpha+1}^*$ given by $f \cup \{(\alpha, i)\}$.⁷

⁷Note that when α is a non-limit ordinal, $\langle i \rangle$ is an element of \mathcal{T}_α . When $f \in \mathcal{T}_\alpha$ and $f \frown \langle i \rangle$ is defined, then $f \frown \langle i \rangle \in \mathcal{T}_\alpha$. Likewise, if $f \in \mathcal{T}_{\alpha+1}$, then $\langle i \rangle \frown f \in \mathcal{T}_\alpha$.

For $\alpha < \beta$, we define the canonical inclusion map $\iota_{\alpha\beta} : \mathcal{T}_\alpha \rightarrow \mathcal{T}_\beta$ by $\iota_{\alpha\beta}(f) = f \cup \{(\gamma, 0) : \gamma \in \beta \setminus \alpha\}$. (Note that $\iota_{\alpha, \alpha+1}(f) = \langle 0 \rangle \frown f$.)

For $\beta < \alpha$, we write ζ_β for the function whose domain is $[\beta, \alpha)$ with the property that $\zeta_\beta(\gamma) = 0$ for all $\gamma \in [\beta, \alpha)$.

Given a family $(b_f)_{f \in X}$, we may refer to it briefly as $b_{\in X}$.

Definition 3.2. We say that a tree $(b_f)_{f \in \mathcal{T}_\alpha}$ is *s-indiscernible over A* if for any tuples $f_0 \dots f_{n-1}$ and $g_0 \dots g_{n-1}$ in \mathcal{T}_α with $f_0 \dots f_{n-1} \equiv^{\text{qf}} g_0 \dots g_{n-1}$, $b_{f_0} \dots b_{f_{n-1}} \equiv_A b_{g_0} \dots b_{g_{n-1}}$, where quantifier-free type is in the language $\mathcal{L}_{s, \alpha}$. (Note that this does not entail that b_f 's on different levels have the same length as tuples.) We also say that $(b_f)_{f \in \mathcal{F}_\alpha}$ is *s-indiscernible over A* (for α a non-limit) if it is *s-indiscernible* after adding a root node $b_{\zeta_\alpha} = \emptyset$.

Given $f \in \mathcal{T}_\alpha$, we write $b_{\geq f}$ to refer to some fixed enumeration of the set $\{b_g : g \in \mathcal{T}_\alpha, f \trianglelefteq g\}$. In particular, we choose this enumeration in a uniform way so that if $(b_f)_{f \in \mathcal{T}_\alpha}$ is *s-indiscernible over A* , then for any f of successor length, the sequence $(b_{f \frown (i)})_{i < \omega}$ is *A-indiscernible*. When f is an element of \mathcal{T}_α^* , we will also write, by an abuse of notation, $b_{\geq f}$ for some fixed enumeration of the set $\{b_g : g \in \mathcal{T}_\alpha, f \subseteq g\}$. One particular example of this will be sequences of the form $(b_{\geq \zeta_{\alpha+1} \frown (i)})_{i < \omega}$, where α is a limit ordinal. This is essentially the only situation in which we need to consider \mathcal{T}_α^* .

Note that for a limit ordinal α , $(b_f)_{f \in \mathcal{T}_\alpha}$ is *s-indiscernible over A* if and only if $(b_f)_{f \in \iota_{\beta, \alpha}(\mathcal{T}_\beta)}$ is *s-indiscernible over A* for every $\beta < \alpha$.

We will also need the following fact.

Fact 3.3 (Modeling property for *s-indiscernibles* [9, Thm. 4.3]). *Let X be \mathcal{T}_α or \mathcal{F}_α . For any $(b_f)_{f \in X}$ and any set A of **hyperimaginaries**, there is a family of tuples $(c_f)_{f \in X}$ that is *s-indiscernible over A* and locally based on $b_{\in X}$ (i.e., for any finite tuple $f_0 \dots f_{n-1}$ from X and any neighborhood U of $\text{tp}(c_{f_0} \dots c_{f_{n-1}}/A)$ (in the appropriate type space), there is a tuple $g_0 \dots g_{n-1}$ from X such that $f_0 \dots f_{n-1} \equiv^{\text{qf}} g_0 \dots g_{n-1}$ and $\text{tp}(b_{g_0} \dots b_{g_{n-1}}) \in U$).*

Note that while Fact 3.3 is normally formulated for discrete logic, the corresponding statement in continuous logic (and therefore also for hyperimaginaries) follows easily from a very soft general argument: Given a metric structure M and a tree $(b_f)_{f \in X}$ of elements of M , pass to a discretization of the theory and apply [9, Thm. 4.3] there. Then take the resulting *s-indiscernible* family back to the original continuous theory. (See [7] for details regarding this kind of construction.)

Now we are ready to prove full existence for \downarrow^{bu} , but we will take the opportunity to prove a certain technical strengthening which we will need later, in the construction of \downarrow^{bu} -Morley trees.

Lemma 3.4. *If $(b_f)_{f \in \mathcal{T}_\alpha}$ is a tree of real elements that is *s-indiscernible over a set of hyperimaginaries A* , then there is a $\gamma > \alpha$ and a tree $(e_f)_{f \in \mathcal{T}_{\gamma+1}}$ such that*

- $e_{\in \mathcal{T}_{\gamma+1}}$ is *s-indiscernible over A* ,
- for each $f \in \mathcal{T}_\alpha$, $b_f = e_{\iota_{\alpha, \gamma+1}(f)}$, and
- $e_{\geq \zeta_\alpha} \downarrow_A^{\text{bu}} e_{\geq h}$, where $\text{dom}(h) = [\alpha, \gamma + 1)$, $h(\delta) = 0$ for all $\delta \in [\alpha, \gamma)$, and $h(\gamma) = 1$.

(Note that $e_{\geq \zeta_\alpha}$ is the original tree.)

Proof. If $b_{\in \mathcal{T}_\alpha} \in \text{acl}(A)$, then the statement is trivial, so assume that $b_{\in \mathcal{T}_\alpha} \notin \text{acl}(A)$.

Fix $\lambda = |Ab_{\in \mathcal{T}_\alpha}| + |T|$. By Proposition 2.3, we have that $b_{\in \mathcal{T}_\alpha} \downarrow_A^{\text{bu}} c$ if and only if $\text{bdd}_\lambda^{\text{u}}(Ab_{\in \mathcal{T}_\alpha}) \cap \text{bdd}_\lambda^{\text{u}}(Ac) = \text{bdd}_\lambda^{\text{u}}(A)$ for any c . Let $\mu = |\text{bdd}_\lambda^{\text{u}}(Ab_{\in \mathcal{T}_\alpha}) \setminus \text{bdd}_\lambda^{\text{u}}(A)|$.

We will build a family $(e_f : f \in \iota_{\gamma+1, \mu^+}(\mathcal{T}_{\gamma+1}))$ inductively, where γ is some successor ordinal less than μ^+ . By an abuse of notation, we will systematically conflate the sets $\iota_{\alpha\mu^+}(\mathcal{T}_\alpha)$ and \mathcal{T}_α (and likewise for $\iota_{\alpha\mu^+}(\mathcal{F}_{\alpha+1})$ and $\mathcal{F}_{\alpha+1}$) for all $\alpha < \mu^+$.⁸ Note that in general this will mean that $e_{\geq \zeta_\beta}$ is the same thing as $e_{\in \mathcal{T}_\beta}$.

Let $e_f = b_f$ for all $f \in \mathcal{T}_\alpha$. Since $b_{\in \mathcal{T}_\alpha} \notin \text{acl}(A)$, we can find a family $(d_f)_{f \in \mathcal{F}_{\alpha+1}}$ extending $e_{\in \mathcal{T}_\alpha}$ such that $(d_{\geq \zeta_{\alpha+1} \frown \langle i \rangle})_{i < \omega}$ is a non-constant A -indiscernible sequence. By Fact 3.3, we can define e_f for all $f \in \mathcal{F}_{\alpha+1}$ in such a way that the family $e_{\in \mathcal{F}_{\alpha+1}}$ is locally based on $d_{\in \mathcal{F}_{\alpha+1}}$. In particular, $(e_{\geq \zeta_{\alpha+1} \frown \langle i \rangle})_{i < \omega}$ will be a non-constant A -indiscernible sequence.

At successor stage $\beta + 1 \geq \alpha$, assume that we have defined e_f for all $f \in \mathcal{F}_{\beta+1}$ and that the family $(e_f)_{f \in \mathcal{F}_{\beta+1}}$ is s -indiscernible over A . If there is no $d_E \in \text{bdd}_\lambda^{\text{u}}(Ab_{\in \mathcal{T}_\alpha}) \setminus \text{bdd}_\lambda^{\text{u}}(A)$ such that the family $(e_f)_{f \in \mathcal{F}_{\beta+1}}$ is s -indiscernible over Ad , let $e_{\zeta_{\beta+1}} = \emptyset$ and $\gamma = \beta$ and halt the construction. Otherwise, let $e_{\zeta_{\beta+1}} = d$. For later reference, let $E_{\beta+1}$ be E . Note that the family $e_{\in \mathcal{T}_{\beta+1}}$ is s -indiscernible over A . Since $d_E \notin \text{bdd}_\lambda^{\text{u}}(A)$, we can find, by Proposition 1.4, a sequence $(\sigma_i)_{i < \omega}$ of elements of $\text{Aut}(\mathbb{M}/A)$ such that for each $i < j < \omega$, $(\sigma_i \cdot d) \not\equiv_{\beta+1} (\sigma_j \cdot d)$. Now chose $(e_f)_{f \in \mathcal{F}_{\beta+2}}$ in such a way that $e_{\in \mathcal{F}_{\beta+2}}$ extends what was already defined, is s -indiscernible over A , and is locally based on the family $(c_f)_{f \in \mathcal{F}_{\beta+2}}$ defined by $c_{\langle i \rangle \frown f} = \sigma_i \cdot e_f$ for all $f \in \mathcal{T}_{\beta+1}$ (which is possible by Fact 3.3). In particular, note that for any $i < j < \omega$, we still have that $(e_{\zeta_{\beta+2} \frown \langle i \rangle}, e_{\zeta_{\beta+2} \frown \langle j \rangle}) \equiv_A (d, \sigma_1 \cdot d)$ and so, in particular, $e_{\zeta_{\beta+2} \frown \langle i \rangle} \not\equiv_{\beta+1} e_{\zeta_{\beta+2} \frown \langle j \rangle}$.

At limit stage β , given $e_{\in \mathcal{T}_\beta}$, note that this family is automatically s -indiscernible over A . Extend it to a family $e_{\in \mathcal{F}_{\beta+1}}$ that is s -indiscernible over A . (This is always possible.)

Claim. For any $\beta < \delta < \mu^+$, if $E_{\beta+1} = E_{\delta+1}$, then $e_{\zeta_{\beta+1}} \not\equiv_{\beta+1} e_{\zeta_{\delta+1}}$.

Proof of claim. The sequence $(e_{\zeta_{\beta+2} \frown \langle i \rangle})_{i < \omega}$ is $e_{\zeta_{\delta+1}}$ -indiscernible. Since

$$e_{\zeta_{\beta+2} \frown \langle 0 \rangle} \not\equiv_{\beta+1} e_{\zeta_{\beta+2} \frown \langle 1 \rangle},$$

it must be the case that $e_{\zeta_{\delta+1}} \not\equiv_{\beta+1} e_{\zeta_{\beta+2} \frown \langle i \rangle}$ for all $i < \omega$. □_{claim}

Let g be the partial function taking β to $[e_{\zeta_{\beta+1}}]_{E_{\beta+1}}$. By the claim, this is an injection into $\text{bdd}_\lambda^{\text{u}}(Ab_{\in \mathcal{T}_\alpha}) \setminus \text{bdd}_\lambda^{\text{u}}(A)$. By the choice of μ , g 's domain cannot be cofinal in μ^+ , so the construction must have halted at some $\gamma < \mu^+$.

Extend $e_{\in \mathcal{T}_\gamma}$ to $e_{\in \mathcal{F}_{\gamma+1}}$ in such a way that the resulting family is s -indiscernible over A . Set $e_{\zeta_{\gamma+1}} = \emptyset$. Let h be the function with domain $[\alpha, \gamma + 1)$ such that $h(\delta) = 0$ for all $\delta \in [\alpha, \gamma)$ and $h(\gamma) = 1$.

Claim. For any $d_E \in \text{bdd}_\lambda^{\text{u}}(Ae_{\geq \zeta_\alpha}) \setminus \text{bdd}_\lambda^{\text{u}}(A)$, $d_E \notin \text{bdd}_\lambda^{\text{u}}(Ae_{\geq h})$.

Proof of claim. Assume that there is some $d_E \in (\text{bdd}_\lambda^{\text{u}}(Ae_{\geq \zeta_\alpha}) \cap \text{bdd}_\lambda^{\text{u}}(Ae_{\geq h})) \setminus \text{bdd}_\lambda^{\text{u}}(A)$. By basic properties of s -indiscernibility, $e_{\in \mathcal{T}_\gamma}$ is s -indiscernible over $Ae_{\geq h}$. By Fact 3.3, we can find a tuple $e' \equiv_A e_{\in \mathcal{T}_\gamma}$ such that $e_{\in \mathcal{F}_{\gamma+1}}$ is s -indiscernible over

⁸Note that the ζ_β and $f \frown \langle i \rangle$ notations are consistent with this conflation but also that the $\langle i \rangle \frown f$ notation is *not*.

Ae' . We can also find a model $M \supseteq Ae'$ with $|M| = \lambda$ such that $e_{\in \mathcal{F}_{\gamma+1}}$ is s -indiscernible over M . By Corollary 1.8, there is an invariant equivalence relation F such that $[M]_F$ and d_E are interdefinable over \emptyset . This implies that $[M]_F \in \text{bdd}_\lambda^u(Ae_{\geq \zeta_\alpha}) \setminus \text{bdd}_\lambda^u(A)$, but this contradicts the fact that the construction halted at step γ . \square_{claim}

So, by the claim, we have that $\text{bdd}_\lambda^u(Ae_{\geq \zeta_\alpha}) \cap \text{bdd}_\lambda^u(Ae_{\geq h}) = \text{bdd}_\lambda^u(A)$. Therefore, by the choice of λ , $e_{\geq \zeta_\alpha} \downarrow_A^{\text{bu}} e_{\geq h}$, as required. \square

Theorem 3.5 (Full existence). *For any set of hyperimaginaries A and real tuples b and c , there is $b' \equiv_A b$ such that $b' \downarrow_A^{\text{bu}} c$.*

Proof. It is sufficient to show this in the special case that $b = c$. Specifically, given d and e , if we can find $d'e' \equiv_A de$ such that $d'e' \downarrow_A^{\text{bu}} de$, then we have $d' \downarrow_A^{\text{bu}} e$ by monotonicity. So fix a set of hyperimaginaries A and a real tuple b . We can now apply Lemma 3.4 to the family $(b_f)_{f \in \mathcal{T}_0}$ with $b_\emptyset = b$ to get a family $(e_f)_{f \in \mathcal{T}_{\gamma+1}}$ such that $e_{\zeta_0} = b$ for some $f \in \mathcal{T}_{\gamma+1}$, $b \equiv_A b_f$ and $b \downarrow_A^{\text{bu}} b_f$. Let σ be an automorphism fixing A taking b_f to b . We then have that $b' = \sigma \cdot b$ is the required element, and we are done. \square

Corollary 3.6. *For any set of hyperimaginaries A and any ultrimaginaries b_E and c_F , there is $b'_E \equiv_A b_E$ such that $b'_E \downarrow_A^{\text{bu}} c_F$.*

Proof. Apply Theorem 3.5 to b and c to get $b' \equiv_A b$ such that $b' \downarrow_A^{\text{bu}} c$. We then have that $\text{bdd}^u(b') \downarrow_A^{\text{bu}} \text{bdd}^u(c)$, so by monotonicity, $b'_E \downarrow_A^{\text{bu}} c_F$. \square

Corollary 3.7 (Extension). *For any set of hyperimaginaries A and any ultrimaginaries b_E , c_F , and d_G , if $b_E \downarrow_A^{\text{bu}} c_F$, then there is $b'_E \equiv_{Ac_F} b_E$ such that $b'_E \downarrow_A^{\text{bu}} c_F d_G$.*

Proof. By Corollary 3.6, we can find $b'_E \equiv_{Ac_F} b$ such that $b'_E \downarrow_{Ac_F}^{\text{bu}} d_G$. By symmetry and transitivity, we have that $b'_E \downarrow_A^{\text{bu}} c_F d_G$. \square

Compactness is very essential in the proof of Fact 3.3 and therefore also Theorem 3.5, which raises the following question.

Question 3.8. *Does Theorem 3.5 hold when A is a set of ultrimaginaries?*

4. TOTAL \downarrow^{bu} -MORLEY SEQUENCES

Definition 4.1. A \downarrow^{bu} -Morley sequence over A is an A -indiscernible sequence $(b_i)_{i < \omega}$ such that $b_i \downarrow_A^{\text{bu}} b_{< i}$ for each $i < \omega$.

A weakly total \downarrow^{bu} -Morley sequence over A is an A -indiscernible sequence $(b_i)_{i < \omega}$ such that for any finite I and any J (of any order type), if $I + J \equiv_A^{\text{EM}} b_{< \omega}$, then $I \downarrow_A^{\text{bu}} J$.

A total \downarrow^{bu} -Morley sequence over A is an A -indiscernible sequence $(b_i)_{i < \omega}$ such that for any I and J (of any order type), if $I + J \equiv_A^{\text{EM}} b_{< \omega}$, then $I \downarrow_A^{\text{bu}} J$.

We could write down stronger and weaker forms of the \downarrow^{bu} -Morley condition, but we are really only interested in total \downarrow^{bu} -Morley sequences, as they seem to be

a fairly robust class (see Theorem 4.8). Weakly total \downarrow^{bu} -Morley sequences seem to be the best we can get without large cardinals, however, which does raise the following question.

Question 4.2. *Is every weakly total \downarrow^{bu} -Morley sequence a total \downarrow^{bu} -Morley sequence?*

One immediate property of total \downarrow^{bu} -Morley sequences is that they act as universal witnesses of Lascar strong type in a strong way.

Proposition 4.3. *For any A and b , if there is a total \downarrow^{bu} -Morley sequence $(b_i)_{i < \omega}$ over A with $b_0 = b$, then for any b' , $b' \equiv_A^L b$ if and only if there are $I_0, J_0, I_1, \dots, J_{n-1}, I_n$ such that $b \in I_0$, $b' \in I_n$, and, for each $i < n$, $I_i + J_i$ and $I_{i+1} + J_i$ are both A -indiscernible and have the same EM-type as $b_{< \omega}$.*

Proof. Let $I = (b_i)_{i < \omega}$. We only need to prove that if $b' \equiv_A^L b$, then the required configuration exists. Choose I' so that $bI \equiv_A^L b'I'$. Then the required configuration exists by Proposition 1.10. \square

A similar statement is true for weakly total \downarrow^{bu} -Morley sequences, which we will state in Corollary 4.17 after we have shown that weakly total \downarrow^{bu} -Morley sequences always exist without set-theoretic hypotheses.

4.1. Characterization of total \downarrow^{bu} -Morley sequences.

Definition 4.4. For any set of parameters A , we write \approx_A for the transitive closure of the relation $I \sim_A J$ that holds if and only if I and J are infinite and either $I + J$ or $J + I$ is an A -indiscernible sequence.

By an abuse of notation, we write $[I]_{\approx_A}$ for the ultraimaginary $[AI]_E$, where E is the equivalence relation on tuples of the same length as AI such that $E(AI, BJ)$ holds if and only if $A = B$ in our fixed enumeration and $I \approx_A J$.

Note that we do not in general require that I and J have the same order type.

We will also need an appropriate Lascar strong type generalization of Ehrenfeucht-Mostowski type.

Definition 4.5. Given two A -indiscernible sequences I and J , we say that I and J have the same *Lascar-Ehrenfeucht-Mostowski type* (or *LEM-type*) over A , written $I \equiv_A^{\text{LEM}} J$, if there is some $J' \equiv_A^L J$ such that $I + J'$ is A -indiscernible.

Lemma 4.6. *For any infinite order types O and O' , $I \approx_A J$ if and only if there are $K_0, L_0, K_1, \dots, L_{n-1}, K_n$ such that*

- $K_0 = I$ and $K_n = J$,
- for $0 < i < n$, K_i is a sequence of order type O ,
- for $i < n$, L_i is a sequence of order type O' , and
- for $i < n$, $K_i + L_i$ and $K_{i+1} + L_i$ are A -indiscernible.

Proof. The \Leftarrow direction is obvious.

For the \Rightarrow direction, we will proceed by induction. First assume that $I \sim_A J$. If $I + J$ is A -indiscernible, then find L of order type O' such that $I + J + L$ is A -indiscernible. We then have that $I + L$ and $J + L$ are A -indiscernible. If $J + I$ is A -indiscernible, then find L of order type O' such that $J + I + L$ is A -indiscernible. We then have that $I + L$ and $J + L$ are A -indiscernible.

Now assume that we know the statement holds for any I and J such that there is a sequence I'_0, \dots, I'_n with $I'_0 = I$, $I'_n = J$, and $I'_i \sim_A I'_{i+1}$ for each $i < n$. Now assume that there is a sequence I'_0, \dots, I'_{n+1} with $I'_0 = I$, $I'_{n+1} = J$, and $I'_i \sim_A I'_{i+1}$ for each $i \leq n$. Apply the induction hypothesis to get $K_0, L_0, K_1, \dots, L_{m-1}, K_m$ satisfying the properties in the statement of the lemma with $K_0 = I$ and $K_m = I'_n$. Now since $I'_n \sim_A I'_{n+1} = J$, we can apply the $n = 1$ case to get L_m such that $I'_n + L_m$ and $I'_{n+1} + L_m$ are both A -indiscernible. By compactness, we can find K_m of order type O such that $K_m + L_m$ and $K_m + L_{m-1}$ are both A -indiscernible. We then have that $K_0, L_0, K_1, \dots, K_m, L_m, K_{m+1}$ is the required sequence, where $K_{m+1} = J$. \square

Proposition 4.7. *Fix a set of hyperimaginary parameters A .*

- (1) \equiv_A^{LEM} is an equivalence relation on the class of A -indiscernible sequences.
- (2) If $I \equiv_A^{\text{L}} J$, then $I \equiv_A^{\text{LEM}} J$.
- (3) If I and J have the same order type, then $I \equiv_A^{\text{L}} J$ if and only if $I \equiv_A^{\text{LEM}} J$.
- (4) If $I \equiv_A^{\text{LEM}} J$, then $I \equiv_A^{\text{EM}} J$.
- (5) If $I \approx_A J$, then $I \equiv_A^{\text{LEM}} J$.

Proof. Recall the following fact: If I and J have the same order type and $I + J$ is A -indiscernible, then $I \equiv_A^{\text{L}} J$.

For (1), it follows easily from the fact that \equiv_A^{LEM} is reflexive. For symmetry, assume that $I \equiv_A^{\text{LEM}} J$, and let I, J , and J' be as in the definition of \equiv_A^{LEM} . Find I' such that $IJ' \equiv_A^{\text{L}} I'J$. Then extend $I' + J$ to $I' + J + I''$, where I'' has the same order type as I . We then have that $I'' \equiv_A^{\text{L}} I' \equiv_A^{\text{L}} I$, so $J \equiv_A^{\text{LEM}} I$. Finally, assume that $I \equiv_A^{\text{LEM}} J$ and $J \equiv_A^{\text{LEM}} K$. Let this be witnessed by J' and K' such that $I + J'$ and $J + K'$ are A -indiscernible. Find K'' with the same order type as K such that $I + J' + K''$ is A -indiscernible. Then find K^* such that $J'K'' \equiv_A^{\text{L}} JK^*$. We have that $K \equiv_A^{\text{L}} K^* \equiv_A^{\text{L}} K''$, so $I \equiv_A^{\text{LEM}} K$.

(2) and (3) are immediate from the fact. (4) is obvious.

For (5), let I and J be such that $I \approx_A J$. Let $K_0, L_0, K_1, \dots, L_{n-1}, K_n$ be as in Lemma 4.6 where the K_i 's and L_i 's (other than K_0) have the same order type as J . We have that $K_i \equiv_A^{\text{L}} L_i$ and $K_i \equiv_A^{\text{L}} L_{i-1}$ for each $i > 0$. Therefore we can take L_0 to be the required J' . \square

Now we will see that total \downarrow^{bu} -Morley sequences over A are precisely those which are 'as generic as possible' in terms of \approx_A (i.e., their \equiv_A^{LEM} -equivalence class decomposes into a single \approx_A -equivalence class).

Theorem 4.8. *For any A -indiscernible sequence $(b_i)_{i < \omega}$ (with A a set of hyperimaginary parameters), the following are equivalent.*

- (1) $b_{< \omega}$ is a total \downarrow^{bu} -Morley sequence over A .
- (2) There exists a pair of infinite sequences I and J (of any, possibly distinct order types) such that $I + J \equiv_A^{\text{EM}} b_{< \omega}$ and $I \downarrow_A^{\text{bu}} J$.
- (3) For any I , $I \approx_A b_{< \omega}$ if and only if $I \equiv_A^{\text{LEM}} b_{< \omega}$.
- (4) $[b_{< \omega}]_{\approx_A} \in \text{bdd}^{\text{u}}(A)$.

Proof. (1) \Rightarrow (2). This is immediate from the definition.

(2) \Rightarrow (3). Let I and J be as in the statement of (2). By Proposition 4.7, we obviously have that if $K \approx_A I$, then $K \equiv_A^{\text{LEM}} I$. Conversely, assume that $K \equiv_A^{\text{LEM}} I$, so there

is $I' \equiv_A^L I$ such that $K + I'$ is A -indiscernible. Since $I \downarrow_A^{\text{bu}} J$, Proposition 2.4 allows us to find $I_0, J_0, I_1, \dots, J_{n-1}, I_n$ such that $I_0 = I$, $J_0 = J$, $I_n = I'$, and, for each $i < n$, $I_i \equiv_{AJ_i}^L I_{i+1}$ and $J_i \equiv_{AJ_{i+1}}^L J_{i+1}$, if $i < n - 1$. In particular, this implies that for each $i < n$, $I_i + J_i$ and $I_{i+1} + J_i$ are both A -indiscernible, so $I \approx_A I' \sim_A K$, whence $I \approx_A K$.

(3) \Rightarrow (1). Since $I \approx_A J$ always implies $I \equiv_A^{\text{LEM}} J$, we only need to prove one direction.

Assume (3), and let I and J be such that $I + J \equiv_A^{\text{EM}} b_{<\omega}$. Let I' be some sequence of the same order type as I . Suppose furthermore that $I \equiv_A^L I'$. This implies that $I \equiv_A^{\text{LEM}} I'$, so, by assumption, we have that $I \approx_A I'$. By Lemma 4.6, we can find $I_0, J_0, I_1, \dots, J_{n-1}, I_n$ such that $I_0 = I$, $J_0 = J$, $I_n = I'$, each I_i has the same order type as I , each J_i has the same order type as J , and $I_i + J_i$ and $I_{i+1} + J_i$ are both A -indiscernible for each $i < n$.

It is generally true that if $K + L$ and $K + L'$ are both A -indiscernible, then $L \equiv_{AK}^L L'$. Therefore, we have that for each $i < n$, $I_i \equiv_{AJ_i}^L I_{i+1}$ and, for each $i < n - 1$, $J_i \equiv_{AJ_{i+1}}^L J_{i+1}$. Since we can do this for any $I' \equiv_A^L I$, Proposition 2.4 yields that $I \downarrow_A^{\text{bu}} J$.

(3) \Rightarrow (4). This follows immediately from Proposition 4.7 and the fact that $[b_{<\omega}] \equiv_A^L$ is always in $\text{bdd}^u(A)$.

(4) \Rightarrow (3). Let $I \equiv_A^{\text{LEM}} b_{<\omega}$. Find I' such that $I \equiv_A^L I'$ and $b_{<\omega} + I'$ is A -indiscernible. Since $[b_{<\omega}] \approx_A \in \text{bdd}^u(A)$, we must also have that $[I'] \approx_A \in \text{bdd}^u(A)$. By Proposition 1.13, there must be an automorphism $\sigma \in \text{Aut}(\mathbb{M}/A, [I'] \approx_A)$ such that $\sigma \cdot I' = I$. Therefore $I \approx_A I'$ and hence $I \approx_A b_{<\omega}$. \square

4.2. Building ((weakly) total) \downarrow^{bu} -Morley sequences. Given that \downarrow^{bu} satisfies full existence, an immediate, familiar Erdős-Rado argument gives that \downarrow^{bu} -Morley sequences exist, but in the end we will need a technical strengthening of this result.

Proposition 4.9. *If $(b_f)_{f \in \mathcal{T}_\alpha}$ is s -indiscernible over A , then there is a family $(c_f)_{f \in \mathcal{F}_{\alpha+1}}$ such that*

- $c_{f \in \mathcal{F}_{\alpha+1}}$ is s -indiscernible over A ,
- $c_{\iota_{\alpha, \alpha+1}(f)} = b_f$ for each $f \in \mathcal{T}_\alpha$, and
- the sequence $(c_{\geq(i)})_{i < \omega}$ is an \downarrow^{bu} -Morley sequence over A .

Proof. Let κ be sufficiently large to apply Erdős-Rado to a sequence of tuples of the same length as $b_{\in \mathcal{T}_\alpha}$ over the set A .

Let $\gamma(0) = \alpha$. Let $c_f^0 = b_f$ for all $f \in \mathcal{T}_{\gamma(0)} = \mathcal{T}_\alpha$. Let $g_0 = \emptyset$ (as an element of \mathcal{T}_α).

At successor stage $\beta + 1$, assume we have $(c_f^\beta)_{\mathcal{T}_{\gamma(\beta)}}$ which is s -indiscernible over A and which satisfies $c_{\iota_{\delta, \beta}(f)}^\beta = c_f^\delta$ for all $\delta \in [\alpha, \beta)$. By Lemma 3.4, we can build a family $(c_f^{\beta+1})_{\mathcal{T}_{\gamma(\beta+1)}}$ (for some successor ordinal $\gamma(\beta + 1) > \gamma(\beta)$) such that

- $(b_f^{\beta+1})_{f \in \mathcal{T}_{\gamma(\beta+1)}}$ is s -indiscernible over A ,
- for each $f \in \mathcal{T}_{\gamma(\beta)}$, $c_f^\beta = c_{\iota_{\gamma(\beta), \gamma(\beta+1)}(f)}^{\beta+1}$, and

- $c_{\geq \zeta(\gamma(\beta))}^{\beta+1} \downarrow_A^{\text{bu}} c_{\geq h}^{\beta+1}$, where $\text{dom}(h) = [\gamma(\beta), \gamma(\beta+1))$, $h(\delta) = 0$ for all $\delta \in [\gamma(\beta), \gamma(\beta+1) - 1)$, and $h(\gamma(\beta+1) - 1) = 1$.

Let $g_{\beta+1} \in \mathcal{T}_{\gamma(\beta+1)}$ be the function with domain $[\alpha, \gamma(\beta+1))$ which has $h(\delta) = 0$ for all $\delta \in [\alpha, \gamma(\beta+1) - 1)$ and $h(\gamma(\beta+1) - 1) = 1$. Note that $g_{\beta+1} \supseteq h$. Also note that by induction we have that

$$c_{\geq g_{\beta+1}}^{\beta+1} \downarrow_A^{\text{bu}} \{c_{\geq \iota_{\gamma(\delta), \gamma(\beta+1)}(g_\delta)}^{\beta+1} : \delta < \beta + 1\},$$

since $\iota_{\gamma(\delta), \gamma(\beta+1)}(g_\delta) \supseteq \zeta_{\gamma(\beta)}$ for all $\delta < \beta + 1$.

At limit stage β , let $\gamma(\beta) = \sup_{\delta < \beta} \gamma(\delta)$ and let $(c_f^\beta)_{f \in \mathcal{T}_{\gamma(\beta)}}$ be the direct limit of $(c_f^\delta)_{f \in \mathcal{T}_{\gamma(\delta)}}$ for $\delta < \beta$. Leave g_β undefined.

Stop once we have $(b_f^\kappa)_{f \in \mathcal{T}_{\gamma(\kappa)}}$. Consider the sequence $(b_{\geq \iota_{\gamma(\beta), \kappa}(g_\beta)}^\kappa)_{\beta \in \kappa \setminus \lim \kappa}$.⁹ By our choice of κ and a standard application of the Erdős-Rado theorem, we can find a family $(c_f)_{f \in \mathcal{F}_{\alpha+1}}$ such that the sequence $(c_{\geq \langle i \rangle})_{i < \omega}$ is A -indiscernible and for every increasing tuple $\bar{i} < \omega$, there is $\bar{\beta} \in \kappa \setminus \lim \kappa$ such that $c_{\geq \langle i_0 \rangle} \cdots c_{\geq \langle i_k \rangle} \equiv_A b_{\geq \iota_{\gamma(\beta_0), \kappa}(g_{\beta_0})}^\kappa \cdots b_{\geq \iota_{\gamma(\beta_k), \kappa}(g_{\beta_k})}^\kappa$.

In particular, note that this implies that

$$c_{\geq \langle i \rangle} \downarrow_A^{\text{bu}} \{c_{\geq \langle j \rangle} : j < i\}$$

for every $i < \omega$. Clearly by applying an automorphism, we may assume that $c_{\iota_{\alpha, \alpha+1}}(f) = b_f$ for each $f \in \mathcal{T}_\alpha$, so all we need to do is show that the family $c_{\in \mathcal{F}_{\alpha+1}}$ is s -indiscernible over A .

Since the sequence $(c_{\geq \langle i \rangle})_{i < \omega}$ is A -indiscernible, it is sufficient, by induction, to show the following statement: For any sequence $\bar{f}_0, \bar{f}_1, \dots, \bar{f}_k, \dots, \bar{f}_\ell$ of tuples of elements of $\mathcal{F}_{\alpha+1}$ satisfying $\bar{f}_i \supseteq \langle i \rangle$ for all $i \leq \ell$ and any $\bar{h} \supseteq \langle k \rangle$ such that \bar{f}_k and \bar{h} realize the same quantifier-free type, we have that $c_{\bar{f}_k}$ and $c_{\bar{h}}$ realize the same type over $Ac_{\bar{f}_0} \cdots c_{\bar{f}_{k-1}} c_{\bar{f}_{k+1}} \cdots c_{\bar{f}_\ell}$.

So let $\bar{f}_0, \dots, \bar{f}_\ell$ and \bar{h} be as in the statement. By construction, there are $\beta_0, \dots, \beta_\ell$ such that $c_{\geq \langle i \rangle} \equiv_A b_{\geq \iota_{\gamma(\beta_i), \kappa}(g_{\beta_i})}^\kappa$ for each $i \leq \ell$. Let $\bar{f}'_0, \dots, \bar{f}'_\ell, \bar{h}'$ be the corresponding elements of \mathcal{T}_κ . (So, in particular, $\bar{f}'_i \supseteq g_{\beta_i}$ for each $i \leq \ell$ and $\bar{h}' \supseteq g_{\beta_k}$). We now have that \bar{f}'_k and \bar{h}' realize the same quantifier-free type. Therefore, by the s -indiscernible of $b_{\in \mathcal{T}_\kappa}^\kappa$, we have that $b_{\bar{f}'_k}^\kappa$ and $b_{\bar{h}'}^\kappa$ realize the same type over $Ab_{\bar{f}'_0}^\kappa \cdots b_{\bar{f}'_{k-1}}^\kappa b_{\bar{f}'_{k+1}}^\kappa \cdots b_{\bar{f}'_\ell}^\kappa$. From this the required statement follows, and we have that $c_{\in \mathcal{F}_{\alpha+1}}$ is s -indiscernible over A . \square

Corollary 4.10. *For any set of hyperimaginaries A and any real tuple b , there is an \downarrow_A^{bu} -Morley sequence $(b_i)_{i < \omega}$ over A with $b_0 = b$.*

Proof. Apply Proposition 4.9 to the tree $(b_f)_{f \in \mathcal{T}_0}$ defined by $b_\emptyset = b$. \square

The order type ω is essential, however; Erdős-Rado only guarantees the existence of sequences that satisfy the relevant condition on finite tuples. Fortunately, this is more than sufficient for the following weak ‘chain condition.’

Lemma 4.11. *If $(b_i)_{i < \omega}$ is an \downarrow_A^{bu} -Morley sequence over A that is moreover Ac -indiscernible, then $b_0 \downarrow_A^{\text{bu}} c$.*

⁹We write $\lim \alpha$ for the set of limit ordinals in α .

Proof. Fix λ . Let $\mu = |\text{bdd}_\lambda^u(Ac) \setminus \text{bdd}_\lambda^u(A)|$. Extend $b_{<\omega}$ to $(b_i)_{i < \mu^+}$. We still have that for any $i < j < \mu^+$, $b_i \downarrow_A^{\text{bu}} b_j$ (since this is only a property of $\text{tp}(b_i b_j / A)$). Therefore the sets $\text{bdd}_\lambda^u(Ab_i) \setminus \text{bdd}_\lambda^u(A)$ are pairwise disjoint. Since there are μ^+ many of them, one of them must be disjoint from $\text{bdd}_\lambda^u(Ac) \setminus \text{bdd}_\lambda^u(A)$. Therefore by indiscernibility, we must have $b_0 \downarrow_A^{\text{bu}} c$. \square

We will not use the following corollary of Lemma 4.11, but it is worth pointing out.

Corollary 4.12. *If I is a total \downarrow^{bu} -Morley sequence over A that is Ac -indiscernible, then $I \downarrow_A^{\text{bu}} c$.*

Proof. Extend I to an Ac -indiscernible sequence $I_0 + I_1 + I_2 + \dots$ with $I_0 = I$. Since I is totally \downarrow^{bu} -Morley, we have that $(I_i)_{i < \omega}$ is an \downarrow^{bu} -Morley sequence over A . So by Lemma 4.11, we have $I = I_0 \downarrow_A^{\text{bu}} c$. \square

Part (2) of following definition is equivalent to [9, Def. 2.1, 3.4] in our context. The rest of it is based on [8, Def. 5.7].

Definition 4.13. Fix a family $(b_f)_{f \in \mathcal{T}_\alpha}$.

- (1) For $w \subseteq \alpha \setminus \lim \alpha$, the restriction of \mathcal{T}_α to the set of levels w is given by $\mathcal{T}_\alpha \upharpoonright w = \{f \in \mathcal{T}_\alpha : \min \text{dom}(f) \in w, \beta \in \text{dom}(f) \setminus w \Rightarrow f(\beta) = 0\}$.
- (2) A family $(b_f)_{f \in \mathcal{T}_\alpha}$ is *str-indiscernible over A* if it is s -indiscernible over A and satisfies that for any $w, v \in [\alpha \setminus \lim \alpha]^{<\omega}$ with $|w| = |v|$, $b_{\in \mathcal{T}_\alpha \upharpoonright w}$ and $b_{\in \mathcal{T}_\alpha \upharpoonright v}$ realize the same type over A .
- (3) We say that $b_{\in \mathcal{T}_\alpha}$ is \downarrow^{bu} -spread out over A if for any $f \in \mathcal{T}_\alpha$ (with $\text{dom}(f) = [\beta + 1, \alpha)$ for some $\beta < \alpha$), the sequence $(b_{\succeq f \frown (i)})_{i < \omega}$ is an \downarrow^{bu} -Morley sequence over A .
- (4) $b_{\in \mathcal{T}_\alpha}$ is an \downarrow^{bu} -Morley tree over A if it is \downarrow^{bu} -spread out and *str-indiscernible over A* .

Note that if $b_{\in \mathcal{T}_\alpha}$ is \downarrow^{bu} -spread out over A , then any restriction $b_{\in \mathcal{T}_\alpha \upharpoonright w}$ is also \downarrow^{bu} -spread out over A (even for infinite w). Also note that, by a basic compactness argument, if α is infinite and $(b_f)_{f \in \mathcal{T}_\alpha}$ is *str-indiscernible over A* , then for any β , we can find a tree $(c_f)_{f \in \mathcal{T}_\beta}$ which is *str-indiscernible over A* such that for any $w \in [\alpha]^{<\omega}$ and $v \in [\beta]^{<\omega}$ with $|w| = |v|$, $b_{\in \mathcal{T}_\alpha \upharpoonright w} \equiv_A c_{\in \mathcal{T}_\beta \upharpoonright v}$.

Proposition 4.14. *For any A, b , and κ , there is a tree $(b_f)_{f \in \mathcal{T}_\kappa}$ that is \downarrow^{bu} -spread out and s -indiscernible over A such that for each $f \in \mathcal{T}_\kappa$, $b_f \equiv_A b$.*

Proof. Let $(b_f^0)_{f \in \mathcal{T}_0}$ be defined by $b_\emptyset^0 = b$. This is vacuously \downarrow^{bu} -spread out and s -indiscernible out over A .

At successor stage $\alpha+1$, given $(b_f^\alpha)_{f \in \mathcal{T}_\alpha}$ which is \downarrow^{bu} -spread out and s -indiscernible by Proposition 4.9, we can find an extension $(b_f^{\alpha+1})_{f \in \mathcal{F}_{\alpha+1}}$ satisfying $b_{\iota_{\alpha, \alpha+1}(f)}^{\alpha+1} = b_f^\alpha$ for all $f \in \mathcal{T}_\alpha$ such that $b_{\in \mathcal{F}_{\alpha+1}}^{\alpha+1}$ is s -indiscernible over A and $(b_{\succeq (i)}^{\alpha+1})_{i < \omega}$ is an \downarrow^{bu} -Morley sequence over A . By Fact 3.3, we can find $b_\emptyset^{\alpha+1} \equiv_A b$ such that the tree $(b_f^{\alpha+1})_{f \in \mathcal{T}_{\alpha+1}}$ is s -indiscernible over A . By construction, we now have that $(b_f^{\alpha+1})_{f \in \mathcal{T}_{\alpha+1}}$ is \downarrow^{bu} -spread out over A .

At limit stage α , let $(b_f^\alpha)_{f \in \mathcal{T}_\alpha}$ be the direct limit of $(b_f^\beta)_{f \in \mathcal{T}_\beta}$ for $\beta < \alpha$. It is immediate from the definitions that $b_{\in \mathcal{T}_\alpha}^\alpha$ is \downarrow^{bu} -spread out and s -indiscernible over A .

Once we have constructed $(b_f^\kappa)_{f \in \mathcal{T}_\kappa}$, let $b_f = b_f^\kappa$ for each $f \in \mathcal{T}_\kappa$. We have that $b_{\in \mathcal{T}_\kappa}$ is the required tree by induction. \square

By the same argument as in [8, Lem. 5.10], we get the following.

Lemma 4.15. *Suppose $(b_f)_{f \in \mathcal{T}_\kappa}$ is \downarrow^{bu} -spread out and s -indiscernible over A with all b_f tuples of the same length. If κ is sufficiently large, then there is an \downarrow^{bu} -Morley tree $(c_f)_{f \in \mathcal{T}_\omega}$ such that for any $w \in [\omega]^{<\omega}$, there is $v \in [\kappa]^{<\omega}$ such that*

$$(b_f)_{f \in \mathcal{T}_\kappa \upharpoonright v} \equiv_A (c_f)_{f \in \mathcal{T}_\omega \upharpoonright w}.$$

Proof. For any tree $(b_f)_{f \in \mathcal{T}_\kappa}$ where all tuples b_f are the same length, define a function t on $[\kappa \setminus \lim \kappa]^{<\omega}$ that takes $w \in [\kappa \setminus \lim \kappa]^{<\omega}$ to $\text{tp}(b_{\in \mathcal{T}_\kappa \upharpoonright w}/A)$. Let κ be large enough to apply Erdős-Rado with the appropriate number of colors.

By Proposition 4.14, we can find a tree $(b_f)_{f \in \mathcal{T}_\kappa}$ that is s -indiscernible and \downarrow^{bu} -spread out over A . By the typical Erdős-Rado argument, we can find a family F of finite sets $w \subset \kappa \setminus \lim \kappa$ closed under subsets such that for any $w, v \in F$ with $|w| = |v|$, $b_{\in \mathcal{T}_\kappa \upharpoonright w} \equiv_A b_{\in \mathcal{T}_\kappa \upharpoonright v}$. By compactness, we can build a tree $(c_f)_{f \in \mathcal{T}_\omega}$ such that for any $w \in [\omega]^{<\omega}$, if $v \in F$ has $|w| = |v|$, then $c_{\in \mathcal{T}_\omega \upharpoonright w} \equiv_A b_{\in \mathcal{T}_\kappa \upharpoonright v}$. Finally, we have that $c_{\in \mathcal{T}_\omega \upharpoonright [0, n]}$ is an \downarrow^{bu} -Morley tree for every $n < \omega$ because $b_{\in \mathcal{T}_\kappa}$ was s -indiscernible and \downarrow^{bu} -spread out over A . Therefore $c_{\in \mathcal{T}_\omega}$ is s -indiscernible and \downarrow^{bu} -spread out over A , and so $c_{\in \mathcal{T}_\omega}$ is an \downarrow^{bu} -Morley tree over A . \square

Proposition 4.16. *If $(b_f)_{f \in \mathcal{T}_\omega}$ is an \downarrow^{bu} -Morley tree over A , then $(b_{\zeta_\beta})_{\beta < \omega}$ is a weakly total \downarrow^{bu} -Morley sequence over A .*

Proof. Fix a linear order O . Let $c_\alpha = b_{\zeta_\alpha}$ for each $\alpha < \omega$.

For each positive $n < \omega$ and each $i < j < \omega$, we have that $b_{\geq \zeta_n \frown (i)} \downarrow_A^{\text{bu}} b_{\geq \zeta_n \frown (j)}$ and that the sequence $(b_{\geq \zeta_n \frown (i)})_{i < \omega}$ is $Ac_{\geq n}$ -indiscernible. By compactness, we can find $(c_i)_{i \in O}$ such that $(c_i)_{i \in \omega + O}$ is A -indiscernible and such that $(b_{\geq \zeta_n \frown (i)})_{i < \omega}$ is $Ac_{\in [n, \omega) + O}$ -indiscernible for each $n < \omega$.

Therefore, by Lemma 4.11, we have that $c_{< n} \downarrow_A^{\text{bu}} c_{\in [n, \omega) + O}$. Hence, $(b_{\zeta_\beta})_{\beta < \omega}$ is a weakly total \downarrow^{bu} -Morley sequence. \square

Corollary 4.17. *For any A and b , there is an A -indiscernible sequence $(b_i)_{i < \omega}$ with $b_0 = b$ such that for any $b' \equiv_A^L b$ and $n < \omega$, there are $I_0, J_0, I_1, J_1, \dots, J_{k-1}, I_k$ with*

- b the first element of I_0 ,
- b' the first element of I_k ,
- $|I_i| = n$ for all $i \leq k$,
- J_i infinite for all $i < k$, and
- $I_i + J_i$ and $I_{i+1} + J_i$ realizing the same EM-type over A as $b_{< \omega}$ for all $i < k$.

We can also arrange it so that I_i is infinite for all $i \leq k$, $|J_i| = n$ for all $i < k$, and $I_i + J_i$ and $I_{i+1} + J_i$ realize the same EM-type over A as $b_{< \omega}$ in the reverse order for all $i < k$ (with the same choice of $b_{< \omega}$ but possibly a different k).

Proof. This follows immediately from the same reasoning as in Proposition 4.3 together with Lemma 4.15 and Proposition 4.16, which establish that weakly total \downarrow^{bu} -Morley sequences always exist. \square

To go further, we will need the following fact from [11]. Recall that the statement $\kappa \rightarrow (\alpha)_\gamma^{<\omega}$ means that whenever $f : [\kappa]^{<\omega} \rightarrow \gamma$ is a function, there is a set $X \subseteq \kappa$ of order type α such that for each $n < \omega$, f is constant on $[X]^n$.

Fact 4.18 (Silver [11]). *For any limit ordinal α , if κ is the smallest cardinal satisfying $\kappa \rightarrow (\alpha)_2^{<\omega}$, then for any $\gamma < \kappa$, $\kappa \rightarrow (\alpha)_\gamma^{<\omega}$. Furthermore, κ is inaccessible.*

The smallest cardinal λ satisfying $\lambda \rightarrow (\alpha)_2^{<\omega}$ is called the Erdős cardinal $\kappa(\alpha)$. In the specific case of $\alpha = \omega$, we will also need the following fact. (See [6, Exercise 7.4.9] for a proof that easily generalizes to a proof of this fact.)

Fact 4.19. *If $\kappa \rightarrow (\omega)_\gamma^{<\omega}$, then $(\gamma^{<\kappa})^+ \rightarrow (\omega + 1)_\gamma^{<\omega}$.*

In particular, if $\kappa(\omega)$ exists, then for any $\gamma < \kappa(\omega)$, $\kappa(\omega)^+ \rightarrow (\omega + 1)_\gamma^{<\omega}$.

Lemma 4.20. *Suppose $(b_f)_{f \in \mathcal{T}_\lambda}$ is \downarrow^{bu} -spread out and s -indiscernible over A with all b_f tuples of the same length. If $\lambda \rightarrow (\omega + 1)_{2^{|Ab|+|T|}}^{<\omega}$, then there is a set $X \subseteq \lambda \setminus \lim \lambda$ with order type $\omega + 1$ such that $b_{\in \mathcal{T}_\lambda \upharpoonright X}$ is an \downarrow^{bu} -Morley tree over A .*

Proof. Let the function t be as in the proof of Lemma 4.15. By assumption, we can find $X \subset \lambda \setminus \lim \lambda$ of order type $\omega + 1$ such that t is homogeneous on X . The tree $b_{\in \mathcal{T}_\lambda \upharpoonright X}$ is clearly s -indiscernible over A , so the only thing to check is that it is \downarrow^{bu} -spread out over A , but this follows from the general fact that restrictions of trees \downarrow^{bu} -spread out over A are \downarrow^{bu} -spread out over A . \square

Theorem 4.21. *For any A and b in any theory T , if there is a cardinal λ satisfying $\lambda \rightarrow (\omega + 1)_{2^{|Ab|+|T|}}^{<\omega}$, then there is a total \downarrow^{bu} -Morley sequence $(b_i)_{i < \omega}$ over A with $b_0 = b$.*

In particular, it is enough if there is an Erdős cardinal $\kappa(\alpha)$ such that $|Ab| + |T| < \kappa(\alpha)$ (for any limit $\alpha \geq \omega$).

Proof. If the Erdős cardinal $\kappa(\alpha)$ exists and $|Ab| + |T| < \kappa(\alpha)$, then by Fact 4.18, we have $2^{|Ab|+|T|} < \kappa(\alpha)$ as well. Then if $\alpha = \omega$, we have that $\kappa(\alpha)^+ \rightarrow (\omega + 1)_{2^{|Ab|+|T|}}^{<\omega}$ by Fact 4.19. If $\alpha > \omega$, we clearly have $\kappa(\alpha) \rightarrow (\omega + 1)_{2^{|Ab|+|T|}}^{<\omega}$ by Fact 4.18. So in any such case we have the required λ .

Let λ be a cardinal such that $\lambda \rightarrow (\omega + 1)_{2^{|Ab|+|T|}}^{<\omega}$ holds. By Proposition 4.14, we can build a tree $(b_f)_{f \in \mathcal{T}_\lambda}$ that is s -indiscernible and \downarrow^{bu} -spread out over A . By Lemma 4.20 and the choice of λ , we can extract an \downarrow^{bu} -Morley tree $(c_f)_{f \in \mathcal{T}_{\omega+1}}$ from this.

By compactness, we can extend this to a tree $(c_f)_{f \in \mathcal{T}_{\omega+\omega}}$ that is *str*-indiscernible over A . We still have that for any $i < j < \omega$,

$$c_{\geq \zeta_{\omega+1} \frown (i)} \downarrow_A^{\text{bu}} c_{\geq \zeta_{\omega+1} \frown (j)}$$

but now we also have that the $(c_{\geq \zeta_{\omega+1} \frown (i)})_{i < \omega}$ is $A \cup \{c_{\zeta_{\omega+i}} : i < \omega\}$ -indiscernible, by *str*-indiscernibility of the full tree $c_{\in \mathcal{T}_{\omega+\omega}}$. Therefore, by Lemma 4.11,

$$c_{\geq \zeta_{\omega+1} \frown (0)} \downarrow_A^{\text{bu}} \{c_{\zeta_{\omega+i}} : i < \omega\},$$

so in particular,

$$\{c_{\zeta_i} : i < \omega\} \downarrow_A^{\text{bu}} \{c_{\zeta_{\omega+i}} : i < \omega\}.$$

Let $d_i = c_{\zeta_{\alpha(i)}}$ for each $i < \omega + \omega$. We have that $(d_i)_{i < \omega + \omega}$ is A -indiscernible. Furthermore, by Theorem 4.8, we have that $d_{<\omega}$ is a total \downarrow^{bu} -Morley sequence. By applying an automorphism, we get the required $b_{<\omega}$. \square

So if we assume that for every λ , there is a κ such that $\kappa \rightarrow (\omega + 1)_\lambda^{<\omega}$, we get that Lascar strong type is always witnessed by total \downarrow^{bu} -Morley sequences in the manner of Proposition 4.3.

The use of large cardinals in Theorem 4.21 leaves an obvious question.

Question 4.22. *Does the statement ‘for every A and b , there is a total \downarrow^{bu} -Morley sequence $(b_i)_{i < \omega}$ over A with $b_0 = b$ ’ have any set-theoretic strength? What if we add cardinality restrictions, such as $|A| + |T| \leq \aleph_0$ and $|b| < \aleph_0$?*

4.3. Total \downarrow^{bu} -Morley sequences in tame theories. Lemma 4.11 can be used to show that \downarrow^{d} implies \downarrow^{bu} (where $b \downarrow_A^{\text{d}} c$ means that $\text{tp}(b/Ac)$ does not divide over A), something which was previously established for bounded hyperimaginary independence, \downarrow^{b} , in [5] and which is folklore for algebraic independence, \downarrow^{a} .

Proposition 4.23. *For any real elements A , b , and c , if $b \downarrow_A^{\text{d}} c$, then $b \downarrow_A^{\text{bu}} c$.*

Proof. Let $(c_i)_{i < \omega}$ be an \downarrow^{bu} -Morley sequence over A with $c_0 = c$. Since $b \downarrow_A^{\text{d}} c$, we may assume that $c_{<\omega}$ is Ab -indiscernible. Therefore, by Lemma 4.11, $b \downarrow_A^{\text{bu}} c$. \square

Corollary 4.24. *If $(b_i)_{i < \omega}$ is a (non-dividing) Morley sequence over A , then it is a total \downarrow^{bu} -Morley sequence over A .* \square

In simple theories, we get the converse.

Proposition 4.25. *Let T be a simple theory. For any A and A -indiscernible sequence I , the following are equivalent.*

- (1) *I is an \downarrow^{f} -Morley sequence over A .*
- (2) *For any J and K with $J+K \equiv_A^{\text{EM}} I$, $J \downarrow_A^{\text{b}} K$ (i.e., $\text{bdd}^{\text{heq}}(AJ) \cap \text{bdd}^{\text{heq}}(AK) = \text{bdd}^{\text{heq}}(A)$).*
- (3) *I is a total \downarrow^{bu} -Morley sequence over A .*

Proof. (1) \Rightarrow (3) is Corollary 4.24. (3) \Rightarrow (2) is obvious.

(2) \Rightarrow (1) follows by the same reasoning as in [1, Sec. 3.1]. Specifically, in simple theories, $b \downarrow_A^{\text{f}} C$ (with $A \subseteq C$) holds if and only if $\text{cb}(\text{stp}(b/C)) \subseteq \text{bdd}^{\text{heq}}(A)$. This implies that \downarrow^{f} satisfies intersection over hyperimaginaries (i.e., if $A_0 \subseteq C \subseteq A_1$, $b \downarrow_{A_0}^{\text{f}} C$, and $b \downarrow_{A_1}^{\text{f}} C$, then $b \downarrow_{\text{bdd}^{\text{heq}}(A_0) \cap \text{bdd}^{\text{heq}}(A_1)}^{\text{f}} C$). By the same argument as in [1, Lem. 3.4], we have that $b \downarrow_A^{\text{f}} C$ if and only if

- (*) there is an AC -indiscernible sequence $(b_i)_{i < \omega}$ with $b_0 = b$ such that for any J and K with $J + K \equiv_A^{\text{EM}} b_{<\omega}$, $J \downarrow_A^{\text{b}} K$.

Assume that (2) holds. Fix $(b_i)_{i < \omega + \omega} \equiv_A^{\text{EM}} I$. $(b_i)_{\omega \leq i < \omega + \omega}$ is $Ab_{<\omega}$ -indiscernible and satisfies (*). Therefore $b_\omega \downarrow_A^{\text{f}} b_{<\omega}$, and we have that $b_{<\omega + \omega}$, and therefore I , is an \downarrow^{f} -Morley sequence over A . \square

On the other hand, there are easy examples in NIP theories (such as DLO) of total \downarrow^{bu} -Morley sequences that are not strict Morley sequences; any sequence generated by an A -invariant type that is not strictly A -invariant will be a total \downarrow^{bu} -Morley sequence over A but not a strict Morley sequence over A . DLO can also be used to show that not every \downarrow^{b} -Morley sequence in a rosy theory is a total \downarrow^{bu} -Morley sequence (e.g., [1, Ex. 3.13]).

In NSOP₁ theories, we get that tree Morley sequences are total \downarrow^{bu} -Morley sequences.

Proposition 4.26. *Let T be an NSOP₁ theory, and let $M \models T$. If I is a tree Morley sequence over M , then it is a total \downarrow^{bu} -Morley sequence over M .*

Proof. Since we are working over a model, Lascar strong types are types. Let J be a sequence realizing the same EM-type as I over M . Find $K \equiv_M I$ such that $K \downarrow_M^K IJ$. Let I' , J' , and K' have the same order type such that $I + I'$, $J + J'$, and $K + K'$ are all M -indiscernible. Since these are tree Morley sequences, we have that $I \downarrow_M^K I'$, $J \downarrow_M^K J'$, and $K \downarrow_M^K K'$. Therefore, by the independence theorem for NSOP₁ theories, we can find I'' and J'' such that $I + I''$, $K + I''$, $K + J''$, and $J + J''$ are all M -indiscernible, so $I \approx_M J$.

Since we can do this for any such J , we have that I is a total \downarrow^{bu} -Morley sequence. \square

The converse is unclear. The argument in the context of simple theories relies on the existence of canonical bases for types.

Question 4.27. *If T is NSOP₁, is every total \downarrow^{bu} -Morley sequence over $M \models T$ a tree Morley sequence over M ?*

REFERENCES

- [1] Hans Adler. Explanation of Independence. *arXiv e-prints*, page math/0511616, November 2005.
- [2] Hans Adler. A geometric introduction to forking and thorn-forking. *Journal of Mathematical Logic*, 09(01):1–20, June 2009.
- [3] Itai Ben Yaacov. Simplicity in compact abstract theories. 03(02):163–191, November 2003.
- [4] Steven Buechler. Canonical bases in some supersimple theories. 1997. Preprint.
- [5] Gabriel Conant and James Hanson. Separation for isometric group actions and hyperimaginary independence. *arXiv e-prints*, page arXiv:2110.07763, October 2021.
- [6] F. R. Drake. *Set theory : an introduction to large cardinals*. North-Holland Pub. Co. American Elsevier Pub. Co, Amsterdam New York, 1974.
- [7] Isaac Goldbring and H. Jerome Keisler. Continuous sentences preserved under reduced products. *arXiv e-prints*, page arXiv:1905.10673, May 2019.
- [8] Itay Kaplan and Nicholas Ramsey. On Kim-independence. *Journal of the European Mathematical Society*, 22(5):1423–1474, January 2020.
- [9] Byunghan Kim, Hyeung-Joon Kim, and Lynn Scow. Tree indiscernibilities, revisited. *Archive for Mathematical Logic*, 53(1-2):211–232, December 2013.
- [10] Saharon Shelah. Simple unstable theories. *Annals of Mathematical Logic*, 19(3):177–203, December 1980.
- [11] Jack Howard Silver. *Some applications of model theory in set theory*. PhD thesis, 1966.
- [12] Pierre Simon. *A Guide to NIP Theories*. Lecture Notes in Logic. Cambridge University Press, 2015.
- [13] Frank O. Wagner. Plus ultra. *Journal of Mathematical Logic*, 15(02):1550008, December 2015.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK, MD 20742, USA
 Email address: jhanson9@umd.edu